

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number  
**WO 01/68055 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 9/14**
- (21) International Application Number: PCT/IB01/00394
- (22) International Filing Date: 16 March 2001 (16.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/189,942 16 March 2000 (16.03.2000) US
- (71) Applicant (*for all designated States except US*): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **HOOVER, Dennis, Jay** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **SHANKER, Ravi, Mysore** [IN/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **FRIESEN, Dwayne, Thomas** [US/US]; 60779 Currant Way, Bend, OR 97702 (US). **LORENZ, Douglas, Alan** [US/US]; 61332 King Jehu Way, Bend, OR 97702 (US). **NIGHTINGALE, James, Alan, Schriver** [US/US]; 62900 Santa Cruz Avenue, Bend, OR 97701 (US).
- (74) Agent: **LUMB, J., Trevor**; c/o Simpson, Alison, Urquhart-Dykes & Lord, 30 Welbeck Street, London W1G 8ER (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- *with international search report*
  - *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 01/68055 A1**

(54) Title: PHARMACEUTICAL COMPOSITIONS OF GLYCOGEN PHOSPHORYLASE INHIBITORS

(57) Abstract: Pharmaceutical compositions comprise a glycogen phosphorylase inhibitor and at least one concentration-enhancing polymer. The composition may be a simple physical mixture of glycogen phosphorylase inhibitor and concentration-enhancing polymer or a dispersion of glycogen phosphorylase inhibitor and polymer.

PHARMACEUTICAL COMPOSITIONS OF GLYCOGEN  
PHOSPHORYLASE INHIBITORS

BACKGROUND OF THE INVENTION

5                   This invention relates to pharmaceutical compositions containing a glycogen phosphorylase inhibitor (GPI) and at least one concentration-enhancing polymer, and the use of such pharmaceutical compositions to treat diabetes, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemias, hyperlipidemia, atherosclerosis and myocardial ischemia in mammals.

10                   In spite of the early discovery of insulin and its subsequent widespread use in the treatment of diabetes, and the later discovery of and use of sulfonylureas (e.g. Chlorpropamide (Pfizer), Glipizide (Pfizer), Tolbutamide (Upjohn), Acetohexamide (E.I. Lilly), Tolazimide (Upjohn)) and biguanides (e.g. Phenformin (Ciba Geigy), Metformin (G. D. Searle)) as oral hypoglycemic agents, the treatment of diabetes

15                   remains less than satisfactory. The use of insulin, necessary in about 10% of diabetic patients in which synthetic hypoglycemic agents are not effective (Type 1 diabetes, insulin dependent diabetes mellitus), requires multiple daily doses, usually by self-injection.

20                   Determination of the proper dosage of insulin requires frequent estimations of the sugar in urine or blood. The administration of an excess dose of insulin causes hypoglycemia, with effects ranging from mild abnormalities in blood glucose to coma, or even death.

25                   Treatment of non-insulin dependent diabetes mellitus (Type 2 diabetes, NIDDM) usually consists of a combination of diet, exercise, oral agents, e.g. sulfonylureas, and in more severe cases, insulin. However, the clinically available hypoglycemics can have

30                   other side effects which limit their use. In any event, where one of these agents may fail in an individual case, another may succeed. A continuing need for hypoglycemic

35

agents, which may have fewer side effects or succeed where others fail, is clearly evident.

Hepatic glucose production is an important target for NIDDM therapy. The liver is the major  
5 regulator of plasma glucose levels in the post absorptive (fasted) state, and the rate of hepatic glucose production in NIDDM patients is significantly elevated relative to normal individuals. Likewise, in the  
10 postprandial (fed) state, where the liver has a proportionately smaller role in the total plasma glucose supply, hepatic glucose production is abnormally high in NIDDM patients.

Glycogenolysis is an important target for interruption of hepatic glucose production. The liver  
15 produces glucose by glycogenolysis (breakdown of the glucose polymer glycogen) and gluconeogenesis (synthesis of glucose from 2- and 3-carbon precursors). Several lines of evidence indicate that glycogenolysis may make an important contribution to hepatic glucose output in  
20 NIDDM. First, in normal post absorptive man, up to 75% of hepatic glucose production is estimated to result from glycogenolysis. Second, patients having liver glycogen storage diseases, including Hers' disease (glycogen phosphorylase deficiency), display episodic hypoglycemia.  
25 These observations suggest that glycogenolysis may be a significant process for hepatic glucose production.

Glycogenolysis is catalyzed in liver, muscle, and brain by tissue-specific isoforms of the enzyme glycogen phosphorylase (GP). This enzyme cleaves the  
30 glycogen macromolecule to release glucose-1-phosphate and a new shortened glycogen macromolecule. Several types of GPIs have been reported to date: glucose and glucose analogs [Martin, J. L. et al., *Biochemistry* 1991, 30, 10101] and caffeine and other purine analogs [Kasvinsky,  
35 P. J. et al., *J. Biol. Chem.* 1978, 253, 3343-3351 and 9102-9106]. These compounds, and GPIs in general, have been postulated to be of potential use for the treatment


of NIDDM by decreasing hepatic glucose production and lowering glycemia [Blundell, T. B. et al. *Diabetologia* 1992, 35, Suppl. 2, 569-576 and Martin et al. *Biochemistry* 1991, 30, 10101].

5                Sites at which GPIs have been reported to bind are the active site, the caffeine or purine binding site, and the ATP or nucleotide binding site. Enzyme activity is also controlled by phosphorylation at a single phosphorylation site, Ser 14. Phosphorylation normally  
10 causes an increase in GP activity due to a conformational change in the GP enzyme. The features of this conformational change have been identified. See, Sprang et al., *Nature* 1988, 336, 215-21. The experimentally determined GP:GPI structures reveal that inhibitor  
15 binding at any of the three binding sites named above reverses the conformational change in GP that normally occurs upon phosphorylation causing the GP enzyme to adopt the conformation of the "inactive," unphosphorylated protein.

20                Several GPIs have been described. See, e.g., Kristiansen et al., U.S. Patent No. 5,952,363; Lundgren et al., EP 884 050 A1; Kristiansen et al., WO 98/50359; Bols, WO 97/31901; and Lundgren et al., WO 97/09040. Most of these compounds are cyclic amines with various  
25 substitutents that generally render them relatively hydrophilic with good water solubility and good potential for absorption. These GPIs, being water soluble, would thus be expected to not have solubility-limited absorption.

30                A new binding site has been recently discovered, together with new glycogen phosphorylase inhibitors which bind to this new site. See EP 0978279 A1. As used herein and in the claims, this new binding site shall be referred to as the "indole pocket  
35 binding site." Four different types of GPIs have been identified so far that bind to the indole pocket binding

site: See WO 96/39385, U.S. Patent No. 5,952,322, and EP 846464 A2 which disclose GPIs of the first type; WO 96/39384 and EP 832065 A1 which disclose GPIs of the second type; and U.S. Patent No. 5,998,463 which  
5 discloses GPIs of the third type. A fourth type is disclosed herein. In general, these compounds have in common the structural feature of one or more fused ring systems comprising a six-membered aromatic ring and a nitrogen-containing heterocycle. Such fused ring systems  
10 can be considered an "indole-like group," indole itself having the structure:



15

It is believed that GPIs which contain the indole-like group bind to the indole pocket binding site of the GP enzyme. GPIs that bind to this indole pocket binding site generally are relatively hydrophobic, have poor  
20 water solubility, and poor bioavailability when dosed conventionally in crystalline form.

Accordingly, what is therefore desired is a composition containing a poorly water soluble GPI that increases the GPI concentration in aqueous solution, does  
25 not adversely effect the ability of the GPI to bind to the GP enzyme, improves relative bioavailability, and is pharmaceutically acceptable.


#### BRIEF SUMMARY OF INVENTION

30 The present invention overcomes the aforesaid drawbacks by providing a pharmaceutical composition comprising a glycogen phosphorylase inhibitor and a concentration-enhancing polymer. The GPI binds to a portion or all portions of the following residues of a  
35 glycogen phosphorylase enzyme:

parent secondary		
	<u>structure</u>	<u>residue number</u>
		13-23
	helix $\alpha$ 1	24-37
5	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80
	strand $\beta$ 1	81-86
		87-88
10	strand $\beta$ 2	89-92
		93
	helix $\alpha$ 3	94-102
		103
	helix $\alpha$ 4	104-115
15		116-117
	helix $\alpha$ 5	118-124
		125-128
	strand $\beta$ 3	129-131
		132-133
20	helix $\alpha$ 6	134-150
		151-152
	strand $\beta$ 4	153-160
		161
	strand $\beta$ 4b	162-163
25		164-166
	strand $\beta$ 5	167-171
		172-173
	strand $\beta$ 6	174-178
		179-190
30	strand $\beta$ 7	191-192
		194, 197
	strand $\beta$ 8	198-209
		210-211
	strand $\beta$ 9	212-216
35	strand $\beta$ 10	219-226, 228-232
		233-236


	strand $\beta$ 11	237-239, 241, 243-247
		248-260
	helix $\alpha$ 7	261-276
	strand $\beta$ 11b	277-281
5	reverse turn	282-289
	helix $\alpha$ 8	290-304.

In a second aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer, the GPI having the general structure of Formula I:



Formula I


In a third aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer, the GPI having the general structure of Formula II:



Formula II


In a fourth aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer, the GPI having the general structure of Formula III:

7



Formula III

In a fifth aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer, the GPI having the general structure of Formula IV:



Formula IV

In a sixth aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer, the GPI having a solubility in aqueous solution, in the absence of the polymer, of less than 1.0 mg/mL at any pH of from 1 to 8.

In a seventh aspect of the invention, a pharmaceutical composition comprises a GPI and a concentration-enhancing polymer. The composition provides a maximum concentration of the GPI in a use environment that is 1.25-fold that of a control composition comprising an equivalent amount of the GPI and free from the polymer. As used herein, a "use environment" can be either the *in vivo* environment of the GI tract of an animal, particularly a human, or the *in vitro* environment of a test solution, such as phosphate



buffered saline (PBS) or a Model Fasted Duodenal (MFD) solution.

In an eighth aspect of the invention, a pharmaceutical composition comprises a GPI and a  
5 concentration-enhancing polymer. The composition provides a relative bioavailability that is at least 1.25 relative to a control composition comprising an equivalent amount of the GPI and free from the polymer.

In another aspect of the invention, a method of  
10 treatment of a mammal having an indication due to atherosclerosis, diabetes, diabetes prevention, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hypercholesterolemia, hypertriglycerdemia, hypertension, myocardial ischemia, hyperglycemia,  
15 hyperinsulimemia, hyperlipidemia, insulin resistance, bacterial infection, tissue ischemia, diabetic cardiomyopathy, or tumor growth inhibition comprises the following steps. A composition of a GPI and a concentration-enhancing polymer is formed. The  
20 composition is then administered to the mammal.

The composition may be dosed in a variety of dosage forms, including both initial release and controlled release dosage forms, the latter including both delayed and sustained release forms. The  
25 composition may include blends of polymers, and may further include other polymers that improve the aqueous concentration of the GPI. The composition may further comprise other constituents that improve the stability, wetting, dissolution, tableting, or processing  
30 characteristics of the composition.

The various aspects of the present invention each have one or more of the following advantages. The compositions increase the concentration of GPI in aqueous solution relative to the crystalline form of the GPI.  
35 The compositions also improve relative bioavailability of the GPI. In addition, the compositions enable the use of

poorly water soluble, hydrophobic GPIs without adversely affecting their binding characteristics.

The foregoing and other objectives, features, and advantages of the invention will be more readily understood upon consideration of the following detailed description of the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention provides compositions of GPIs and at least one concentration-enhancing polymer. As discussed above in the Background, a new class of poorly water soluble, hydrophobic GPIs has been discovered that bind to the indole pocket binding site in the GP enzyme. It is believed that an important part of the binding of GPIs to this site is due to the indole-like group, which, being relatively hydrophobic, binds in a hydrophobic pocket within the GP enzyme. In studying the GPI activity, binding mode, and GPI/GP complex structure of a wide variety of compounds, it has been found that compounds that have good GP inhibition activity at this indole pocket binding site often have a number of features in common: (1) the presence of one or more indole-like groups in the structure; (2) extremely low solubilities in aqueous solution (*i.e.*, less than 1.0 mg/mL) at physiologically relevant pH (*e.g.*, any pH of from 1 through 8) measured at about 22°C; (3) a relatively hydrophobic nature; and (4) a relatively low bioavailability when orally dosed in the crystalline state.

Accordingly, unlike other previously known GPIs, GPIs which bind to the indole pocket binding site typically require some kind of modification or formulation to enhance their solubility and thereby achieve good bioavailability. However, the inventors have found that many of the conventional methods used to improve solubility, and in turn bioavailability, have proved problematic. One method used generally to improve

drug bioavailability is to form an ionic form of the drug, typically by incorporating an ionizable group into its structure, and particularly by forming a highly soluble salt form. However, the GPIs with the indole-like group having the best performance generally are neutral or nonionic and relatively hydrophobic.

The inventors have found that preparing GPIs having indole-like groups as compositions comprising a GPI and concentration-enhancing polymer, and preferably as a solid dispersion of the GPI and concentration-enhancing polymer, improves the aqueous concentration of the GPIs as well as relative bioavailability, but does not adversely affect the binding characteristics of the GPIs. The compositions, GPIs, suitable polymers, and optional excipients are discussed in more detail as follows.

#### COMPOSITIONS OF GPIs AND CONCENTRATION-ENHANCING POLYMER

The present invention finds utility with any low-solubility GPI, or any GPI which would benefit by improved bioavailability. The compositions of the present invention are mixtures comprised of a GPI and at least one concentration-enhancing polymer. The mixtures are preferably solid dispersions, but simple physical mixtures of the GPI and polymer may also be suitable for some GPIs. The GPI in its pure state may be crystalline or amorphous. Preferably, at least a major portion of the GPI in the composition is amorphous. By "amorphous" is meant simply that the GPI is in a non-crystalline state. As used herein, the term "a major portion" of the GPI means that at least 60% of the GPI in the composition is in the amorphous form, rather than the crystalline form. Preferably, the GPI in the composition is substantially amorphous. As used herein, "substantially amorphous" means that the amount of the GPI in crystalline form does not exceed 25%. More preferably, the GPI in the composition is "almost completely

amorphous" meaning that the amount of GPI in the crystalline form does not exceed 10%. Amounts of crystalline GPI may be measured by powder X-ray diffraction, Scanning Electron Microscope (SEM) analysis, differential scanning calorimetry ("DSC"), or any other standard quantitative measurement. The composition may contain from about 1 to about 80 wt% GPI, depending on the dose of the GPI. Enhancement of aqueous GPI concentrations and relative bioavailability are typically best at low GPI levels, typically less than about 25 to 40 wt%. However, due to the practical limit of the dosage form size, higher GPI loadings are often preferred and perform well.

In a preferred aspect of the invention, GPI and concentration-enhancing polymer are present as a solid dispersion of the low-solubility GPI and polymer. Preferably, at least a major portion of the GPI in the dispersion is present in the amorphous, rather than the crystalline state. The amorphous GPI can exist as a pure phase, as a solid solution of GPI homogeneously distributed throughout the polymer or any combination of these states or those states that lie intermediate between them.

The dispersion is preferably substantially homogeneous so that the amorphous GPI is dispersed as homogeneously as possible throughout the polymer. As used herein, "substantially homogeneous" means that the GPI present in relatively pure amorphous domains within the solid dispersion is relatively small, on the order of less than 20%, and preferably less than 10% of the total amount of GPI. While the dispersion may have some GPI-rich domains, it is preferred that the dispersion itself have a single glass transition temperature ( $T_g$ ) which demonstrates that the dispersion is substantially homogeneous. This contrasts with a simple physical mixture of pure amorphous GPI particles and pure amorphous polymer particles which generally display two

distinct  $T_g$ s, one that of the GPI and one that of the polymer.  $T_g$  as used herein is the characteristic temperature where a glassy material, upon gradual heating, undergoes a relatively rapid (e.g., 10 to 100  
5 seconds) physical change from a glass state to a rubber state. Dispersions of the present invention that are substantially homogeneous generally are more physically stable and have improved concentration-enhancing properties and, in turn improved bioavailability,  
10 relative to nonhomogeneous dispersions.

While the inventors have found that dispersions of the GPI and concentration-enhancing polymer yield good results, it has been found for at least one GPI that compositions of physical mixtures of amorphous GPI and  
15 concentration-enhancing polymer also yield improved aqueous GPI concentration. At least a major portion of the GPI in the mixture is amorphous. The composition may be in the form of a simple dry physical mixture wherein both the GPI and concentration-enhancing polymer are  
20 mixed in particulate form and wherein the particles of each, regardless of size, retain the same individual physical properties that they exhibit in bulk. Any conventional method used to mix the polymer and GPI together such as physical mixing and dry or wet  
25 granulation may be used. In this embodiment of the invention, the amorphous GPI and concentration-enhancing polymer need not be directly mixed, but only both present in the dosage form. For example, the amorphous GPI may be in the form of a tablet, bead, or capsule, and the  
30 concentration-enhancing polymer may be a coating, granulating material, or even the wall of the capsule.

The compositions comprising the GPI and concentration-enhancing polymer provide enhanced concentration of the GPI in *in vitro* dissolution tests.  
35 It has been determined that enhanced drug concentration in *in vitro* dissolution tests in Model Fasted Duodenal (MFD solution) or Phosphate Buffered Saline (PBS) is a

good indicator of *in vivo* performance and bioavailability. An appropriate PBS solution is an aqueous solution comprising 20 mM sodium phosphate ( $\text{Na}_2\text{HPO}_4$ ), 47 mM potassium phosphate ( $\text{KH}_2\text{PO}_4$ ), 87 mM NaCl, and 0.2 mM KCl, adjusted to pH 6.5 with NaOH. An appropriate MFD solution is the same PBS solution wherein additionally is present 14.7 mM sodium taurocholic acid and 2.8 mM of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine. In particular, a composition of the present invention can be dissolution-tested by adding it to MFD or PBS solution and agitating to promote dissolution. Preferably, the composition of the present invention provides a Maximum Drug Concentration (MDC) that is at least 1.25-fold the equilibrium concentration of a control composition comprising an equivalent quantity of GPI but free from the polymer. In other words, if the equilibrium concentration provided by the control composition is 100  $\mu\text{g/mL}$ , then a composition of the present invention provides an MDC of at least 125  $\mu\text{g/mL}$ . The comparison composition is conventionally the undispersed GPI alone (e.g., typically, the crystalline GPI alone in its most thermodynamically stable crystalline form, or in cases where a crystalline form of the GPI is unknown, the control may be the amorphous GPI alone) or the GPI plus a weight of inert diluent equivalent to the weight of polymer in the test composition. More preferably, the MDC of GPI achieved with the compositions of the present invention are at least 2-fold, and even more preferably at least 3-fold, that of the control composition.

Alternatively, the compositions of the present invention provide in an aqueous use environment a concentration versus time Area Under The Curve (AUC), for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment

that is at least 1.25-fold that of a control composition comprising an equivalent quantity of undispersed GPI.

Alternatively, the dispersion of the present invention, when dosed orally to a human or other animal, provides an AUC in GPI concentration in the blood for any period of at least 90 minutes between the time of dosage and about 270 minutes following dosage that is at least 1.25-fold that observed when a control composition comprising an equivalent quantity of undispersed drug is dosed. Thus, the compositions of the present invention can be evaluated in either an *in vitro* or *in vivo* test, or both.

A typical test to evaluate enhanced drug concentration can be conducted by (1) dissolving a sufficient quantity of control composition, typically the GPI alone, in the *in vitro* test medium, typically MFD or PBS solution, to achieve equilibrium concentration of the GPI; (2) dissolving a sufficient quantity of test composition (e.g., the GPI and polymer) in an equivalent test medium, such that if all the GPI dissolved, the theoretical concentration of GPI would exceed the equilibrium concentration of the GPI by a factor of at least 2; and (3) determining whether the measured MDC of the test composition in the test medium is at least 1.25-fold that of the equilibrium concentration of the control composition. In conducting such a dissolution test, the amount of test composition or control composition used is an amount such that if all of the GPI dissolved the GPI concentration would be at least 2-fold to 100-fold that of the solubility of the GPI. The concentration of dissolved GPI is typically measured as a function of time by sampling the test medium and plotting GPI concentration in the test medium vs. time so that the MDC can be ascertained. To avoid GPI particulates which would give an erroneous determination, the test solution is either filtered or centrifuged. "Dissolved GPI" is typically taken as that material that either passes a

0.45  $\mu\text{m}$  syringe filter or, alternatively, the material that remains in the supernatant following centrifugation. Filtration can be conducted using a 13 mm, 0.45  $\mu\text{m}$  polyvinylidene difluoride syringe filter sold by Scientific Resources under the trademark TITAN®.

Centrifugation is typically carried out in a polypropylene microcentrifuge tube by centrifuging at 13,000 G for 60 seconds. Other similar filtration or centrifugation methods can be employed and useful results obtained. For example, using other types of microfilters may yield values somewhat higher or lower ( $\pm 10\text{-}40\%$ ) than that obtained with the filter specified above but will still allow identification of preferred dispersions. It is recognized that this definition of "dissolved GPI" encompasses not only monomeric solvated GPI molecules but also a wide range of species such as polymer/GPI assemblies that have submicron dimensions such as GPI aggregates, aggregates of mixtures of polymer and GPI, micelles, polymeric micelles, colloidal particles or nanocrystals, polymer/GPI complexes, and other such GPI-containing species that are present in the filtrate or supernatant in the specified dissolution test.

Relative bioavailability of GPIs in the dispersions of the present invention can be tested *in vivo* in animals or humans using conventional methods for making such a determination. An *in vivo* test, such as a crossover study, may be used to determine whether a composition of GPI and polymer provides an enhanced relative bioavailability compared with a control composition comprised of a GPI but no polymer as described above. In an *in vivo* crossover study a "test composition" of GPI and polymer is dosed to half a group of test subjects and, after an appropriate washout period (e.g., one week) the same subjects are dosed with a "control composition" that comprises an equivalent quantity of GPI as the "test composition". The other



half of the group is dosed with the control composition first, followed by the test composition. The relative bioavailability is measured as the concentration in the blood (serum or plasma) versus time area under the curve (AUC) determined for the test group divided by the AUC in the blood provided by the control composition. Preferably, this test/control ratio is determined for each subject, and then the ratios are averaged over all subjects in the study. *In vivo* determinations of AUC can be made by plotting the serum or plasma concentration of drug along the ordinate (y-axis) against time along the abscissa (x-axis). Generally, the values for AUC represent a number of values taken from all of the subjects in a patient test population averaged over the entire test population. A preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 1.25 relative to a control composition comprised of a GPI but with no polymer as described above. (That is, the AUC provided by the test composition is at least 1.25-fold the AUC provided by the control composition.) An even more preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 2.0 relative to a control composition of the GPI but with no polymer present, as described above. The determination of AUCs is a well-known procedure and is described, for example, in Welling, "Pharmacokinetics Processes and Mathematics," ACS Monograph 185 (1986).

### 30 GLYCOGEN PHOSPHORYLASE INHIBITORS

The invention is useful for GPIs which have sufficiently low aqueous solubility that it is desirable to increase their water solubility. Therefore, anytime one finds it desirable to raise the concentration of the GPI in a use environment, the invention will find utility. The GPI has "low-solubility," meaning that the GPI may be either "substantially water-insoluble" (which

means that the GPI has a minimum aqueous solubility at any physiologically relevant pH (e.g., pH 1-8) and about 22°C of less than 0.01 mg/mL), or "sparingly water-soluble" (that is, has a water solubility up to about 1 mg/mL). (Unless otherwise specified, reference to aqueous solubility herein and in the claims is determined at about 22°C.) Compositions of the present invention find greater utility as the solubility of the GPI decreases, and thus are preferred for GPI solubilities less than 0.5 mg/mL, and even more preferred for GPI solubilities less than 0.1 mg/mL. In general, it may be said that the GPI has a dose-to-aqueous solubility ratio greater than about 10 mL, where the solubility (mg/mL) is the minimum value observed in any physiologically relevant aqueous solution (e.g., those with pH values from 1 to 8) including USP simulated gastric and intestinal buffers, and dose is in mg. Compositions of the present invention, as mentioned above, find greater utility as the solubility of the GPI decreases and the dose increases. Thus, the compositions are preferred as the dose-to-solubility ratio increases, and thus are preferred for dose-to-solubility ratios greater than 100 mL, and more preferred for dose-to-solubility ratios greater than 400 mL.

Preferably, the GPI binds to the GP enzyme at the indole pocket binding site. As used herein and in the claims, "bind" means a portion of the GPI binds to the GP enzyme in such a manner that a portion of the GPI is in van der Waals or hydrogen bonding contact with a portion or all portions of certain residues of the binding site. In a preferred embodiment, the GPI binds to the GP enzyme with a portion or all portions of the following residues of GP:

18

	parent secondary <u>structure</u>	<u>residue number</u>
		13-23
5	helix $\alpha$ 1	24-37
	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80
	strand $\beta$ 1	81-86
10		87-88
	strand $\beta$ 2	89-92
		93
	helix $\alpha$ 3	94-102
		103
15	helix $\alpha$ 4	104-115
		116-117
	helix $\alpha$ 5	118-124
		125-128
	strand $\beta$ 3	129-131
20		132-133
	helix $\alpha$ 6	134-150
		151-152
	strand $\beta$ 4	153-160
		161
25	strand $\beta$ 4b	162-163
		164-166
	strand $\beta$ 5	167-171
		172-173
	strand $\beta$ 6	174-178
30		179-190
	strand $\beta$ 7	191-192
		194, 197
	strand $\beta$ 8	198-209
		210-211
35	strand $\beta$ 9	212-216
	strand $\beta$ 10	219-226, 228-232
		233-236

19

	strand $\beta$ 11	237-239, 241, 243-247 248-260
	helix $\alpha$ 7	261-276
	strand $\beta$ 11b	277-281
5	reverse turn	282-289
	helix $\alpha$ 8	290-304

More preferably, the GPI binds with one or more of the following residues of GP in one or both subunits:

10	parent secondary <u>structure</u>	<u>residue number</u>
		13-23
15	helix $\alpha$ 1	24-37
	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78 79-80
	strand $\beta$ 2	91-92
20		93
	helix $\alpha$ 3	94-102 103
	helix $\alpha$ 4	104-115 116-117
25	helix $\alpha$ 5	118-124 125-128
	strand $\beta$ 3	129-130
	strand $\beta$ 4	159-160 161
30	strand $\beta$ 4b	162-163 164-166
	strand $\beta$ 5	167-168
	strand $\beta$ 6	178 179-190
35	strand $\beta$ 7	191-192 194, 197
	strand $\beta$ 9	198-200
	strand $\beta$ 10	220-226 228-232

20

		233-236
	strand $\beta$ 11	237-239, 241, 243-247
		248-260
	helix $\alpha$ 7	261-276
5	strand $\beta$ 11b	277-280

Even more preferably, the GPI binds with one or more of the following residues of GP in one or both subunits:

10	<u>residue number</u>
	33-39
	49-66
	94
	98
15	102
	125-126
	160
	162
	182-192
20	197
	224-226
	228-231
	238-239
	241
25	245
	247

Most preferably, the GPI binds with one or more of the following residues of GP in one or both subunits:


30	<u>residue number</u>
	37-39
	53
	57
35	60
	63-64
	184-192

226

229

The indole pocket binding site is disclosed more fully in commonly assigned U.S. provisional patent application Serial No. 95790 filed August 7, 1998 and corresponding published European Patent Application No. EP0978279 A1, the relevant disclosure of which is herein incorporated by reference.

It is believed that certain compounds are capable of binding at the indole pocket binding site. Accordingly, preferred GPIs of the present invention are those that are capable of binding at this site. One such set of compounds has the structure of Formula I:



Formula I

and the pharmaceutically acceptable salts and prodrugs thereof wherein the dotted line (---) is an optional bond, and the various substituents of Formula I are as follows;

A is  $-C(H)=$ ,  $-C((C_1-C_4)\text{alkyl})=$  or  $-C(\text{halo})=$  when the dotted line (---) is a bond, or A is methylene or  $-CH((C_1-C_4)\text{alkyl})-$  when the dotted line (---) is not a bond;

$R_1$ ,  $R_{10}$  or  $R_{11}$  are each independently H, halo, 4-, 6- or 7-nitro, cyano,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, fluoromethyl, difluoromethyl or trifluoromethyl;

$R_2$  is H;

$R_3$  is H or  $(C_1-C_5)$ alkyl;

$R_4$  is methyl, ethyl, n-propyl, hydroxy $(C_1-C_3)$ alkyl,  $(C_1-C_3)$ alkoxy $(C_1-C_3)$ alkyl, phenyl $(C_1-C_4)$ alkyl, phenylhydroxy $(C_1-C_4)$ alkyl, phenyl $(C_1-C_4)$ alkoxy $(C_1-C_4)$ alkyl, thien-2- or -3-yl $(C_1-C_4)$ alkyl or fur-2- or -3-yl $(C_1-C_4)$ alkyl wherein said  $R_4$  rings are mono-, di- or tri-substituted independently on carbon with H, halo,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, trifluoromethyl, hydroxy, amino or cyano;

10 or

$R_4$  is pyrid-2-, -3- or -4-yl $(C_1-C_4)$ alkyl, thiazol-2-, -4- or -5-yl $(C_1-C_4)$ alkyl, imidazol-1-, -2-, -4- or -5-yl $(C_1-C_4)$ alkyl, pyrrol-2- or -3-yl $(C_1-C_4)$ alkyl, oxazol-2-, -4- or -5-yl $(C_1-C_4)$ alkyl, pyrazol-3-, -4- or -5-yl $(C_1-C_4)$ alkyl, isoxazol-3-, -4-, -5-yl $(C_1-C_4)$ alkyl, isothiazol-3-, -4-, -5-yl $(C_1-C_4)$ alkyl, pyridazin-3- or -4-yl $(C_1-C_4)$ alkyl, pyrimidin-2-, -4-, -5- or -6-yl $(C_1-C_4)$ alkyl, pyrazin-2- or -3-yl $(C_1-C_4)$ alkyl or 1,3,5-triazin-2-yl $(C_1-C_4)$ alkyl, wherein said preceding  $R_4$  heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

15

20

$R_5$  is H, hydroxy, fluoro,  $(C_1-C_5)$ alkyl,  $(C_1-C_5)$ alkoxy,  $(C_1-C_6)$ alkanoyl, amino  $(C_1-C_4)$  alkoxy, mono-N- or di-N,N- $(C_1-C_4)$ alkylamino $(C_1-C_4)$ alkoxy, carboxy $(C_1-C_4)$ alkoxy,  $(C_1-C_5)$ alkoxy-carbonyl $(C_1-C_4)$ alkoxy, benzyloxycarbonyl $(C_1-C_4)$ alkoxy, or carbonyloxy wherein said carbonyloxy is carbon-carbon linked with phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding  $R_5$  rings are optionally mono-substituted with halo,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, hydroxy, amino or trifluoromethyl and said mono-substituents are bonded to carbon;

25

30

35

$R_7$  is H, fluoro or  $(C_1-C_5)$ alkyl; or

$R_5$  and  $R_7$  can be taken together to be oxo;  
 $R_6$  is carboxy,  $(C_1-C_8)$ alkoxycarbonyl,  $C(O)NR_8R_9$   
or  $C(O)R_{12}$  wherein

$R_8$  is  $(C_1-C_3)$ alkyl, hydroxy or  $(C_1-C_3)$ alkoxy; and  
5  $R_9$  is H,  $(C_1-C_8)$ alkyl, hydroxy,  $(C_1-C_8)$ alkoxy,  
methylene-perfluorinated  $(C_1-C_8)$ alkyl, phenyl, pyridyl,  
thienyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl,  
thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl,  
pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl,  
10 piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl,  
pyrazinyl, piperazinyl or 1,3,5-triazinyl wherein said  
preceding  $R_9$  rings are carbon-nitrogen linked; or  
 $R_9$  is mono-, di- or tri-substituted  
 $(C_1-C_5)$ alkyl, wherein said substituents are independently  
15 H, hydroxy, amino, mono-N- or di-N,N-  $(C_1-C_5)$ alkylamino; or  
 $R_9$  is mono- or di-substituted  $(C_1-C_5)$ alkyl,  
wherein said substituents are independently phenyl,  
pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl,  
thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl,  
20 pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl,  
pyridinyl, piperidinyl, morpholinyl, pyridazinyl,  
pyrimidinyl, pyrazinyl, piperazinyl or 1,3,5-triazinyl  
wherein the nonaromatic nitrogen-containing  $R_9$   
rings are optionally mono-substituted on nitrogen with  
25  $(C_1-C_6)$ alkyl, benzyl, benzoyl or  $(C_1-C_6)$ alkoxycarbonyl and  
wherein the  $R_9$  rings are optionally mono-substituted on  
carbon with halo,  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, hydroxy,  
amino, or mono-N- and di-N,N-  $(C_1-C_5)$ alkylamino provided  
that no quaternized nitrogen is included and there are no  
30 nitrogen-oxygen, nitrogen-nitrogen or nitrogen-halo  
bonds;

$R_{12}$  is piperazin-1-yl, 4- $(C_1-C_4)$ alkylpiperazin-1-  
yl, 4-formylpiperazin-1-yl, morpholino, thiomorpholino,  
1-oxothiomorpholino, 1,1-dioxo-thiomorpholino,  
35 thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-  
thiazolidin-3-yl, 2- $(C_1-C_6)$ alkoxycarbonylpyrrolidin-1-yl,  
oxazolidin-3-yl or 2(R)-hydroxymethylpyrrolidin-1-yl; or



$R_{12}$  is 3- and/or 4-mono- or di-substituted oxazetidin-2-yl, 2-, 4-, and/or 5-mono- or di-substituted oxazolidin-3-yl, 2-, 4-, and/or 5-mono- or di-substituted thiazolidin-3-yl, 2-, 4- and/or 5-mono- or di-substituted 1-oxothiazolidin-3-yl, 2-, 4-, and/or 5-mono- or di-substituted 1,1-dioxothiazolidin-3-yl, 3- and/or 4-, mono- or di-substituted pyrrolidin-1-yl, 3-, 4- and/or 5-, mono-, di- or tri-substituted piperidin-1-yl, 3-, 4-, and/or 5-mono-, di-, or tri-substituted piperazin-1-yl, 3-substituted azetidin-1-yl, 4- and/or 5-, mono- or di-substituted 1,2-oxazinan-2-yl, 3-and/or 4-mono- or di-substituted pyrazolidin-1-yl, 4- and/or 5-, mono- or di-substituted isothiazolidin-2-yl, 4- and/or 5-, mono- and/or di-substituted isothiazolidin-2-yl wherein said  $R_{12}$  substituents are independently H, halo,  $(C_1-C_5)$ alkyl, hydroxy, amino, mono-N- or di-N,N- $(C_1-C_5)$ alkylamino, formyl, oxo, hydroxyimino,  $(C_1-C_5)$ alkoxy, carboxy, carbamoyl, mono-N- or di-N,N- $(C_1-C_4)$ alkylcarbamoyl,  $(C_1-C_4)$ alkoxyimino,  $(C_1-C_4)$ alkoxymethoxy,  $(C_1-C_6)$ alkoxycarbonyl, carboxy $(C_1-C_5)$ alkyl or hydroxy $(C_1-C_5)$ alkyl;


with the proviso that if  $R_4$  is H, methyl, ethyl or n-propyl,  $R_5$  is OH;

with the proviso that if  $R_5$  and  $R_7$  are H, then  $R_4$  is not H, methyl, ethyl, n-propyl, hydroxy $(C_1-C_3)$ alkyl or  $(C_1-C_3)$ alkoxy $(C_1-C_3)$ alkyl and  $R_6$  is  $C(O)NR_8R_9$ ,  $C(O)R_{12}$  or  $(C_1-C_4)$ alkoxycarbonyl.

Compounds of Formula I are disclosed in published Patent Cooperation Treaty Application number WO 96/39385, the complete disclosure of which is hereby incorporated by reference.

In yet another preferred aspect of the invention, the GPI has the structure of Formula II, which is another class of compounds thought capable of binding to the indole pocket binding site:

25



Formula II

and the pharmaceutically acceptable salts and prodrugs thereof wherein the dotted line (---) is an optional bond and the substituents of Formula II are as follows:

A is  $-C(H)=$ ,  $-C((C_1-C_4)\text{alkyl})=$ ,  $-C(\text{halo})=$  or  $-N=$ , when the dotted line (---) is a bond, or A is methylene or  $-CH((C_1-C_4)\text{alkyl})-$ , when the dotted line (---) is not a bond;

$R_1$ ,  $R_{10}$  or  $R_{11}$  are each independently H, halo, cyano, 4-, 6- or 7-nitro,  $(C_1-C_4)\text{alkyl}$ ,  $(C_1-C_4)\text{alkoxy}$ , fluoromethyl, difluoromethyl or trifluoromethyl;

$R_2$  is H;

$R_3$  is H or  $(C_1-C_5)\text{alkyl}$ ;

$R_4$  is H, methyl, ethyl, n-propyl, hydroxy $(C_1-C_3)\text{alkyl}$ ,  $(C_1-C_3)\text{alkoxy}(C_1-C_3)\text{alkyl}$ , phenyl $(C_1-C_4)\text{alkyl}$ , phenylhydroxy $(C_1-C_4)\text{alkyl}$ , (phenyl) $((C_1-C_4)\text{-alkoxy})(C_1-C_4)\text{alkyl}$ , thien-2- or -3-yl $(C_1-C_4)\text{alkyl}$  or fur-2- or -3-yl $(C_1-C_4)\text{alkyl}$  wherein said  $R_4$  rings are mono-, di- or tri-substituted independently on carbon with H, halo,  $(C_1-C_4)\text{alkyl}$ ,  $(C_1-C_4)\text{alkoxy}$ , trifluoromethyl, hydroxy, amino, cyano or 4,5-dihydro-1H-imidazol-2-yl; or

$R_4$  is pyrid-2-, -3- or -4-yl $(C_1-C_4)\text{alkyl}$ , thiazol-2-, -4- or -5-yl $(C_1-C_4)\text{alkyl}$ , imidazol-2-, -4-, or -5-yl $(C_1-C_4)\text{alkyl}$ , pyrrol-2- or -3-yl $(C_1-C_4)\text{alkyl}$ , oxazol-2-, -4- or -5-yl $(C_1-C_4)\text{alkyl}$ , pyrazol-3-, -4- or -5-yl $(C_1-C_4)\text{alkyl}$ , isoxazol-3-, -4- or -5-yl $(C_1-C_4)\text{alkyl}$ , isothiazol-3-, -4- or -5-yl $(C_1-C_4)\text{alkyl}$ , pyridazin-3- or -4-yl $(C_1-C_4)\text{alkyl}$ , pyrimidin-2-, -4-, -5- or -6-yl $(C_1-C_4)\text{alkyl}$ , pyrazin-2- or -3-yl $(C_1-C_4)\text{alkyl}$ ,

1,3,5-triazin-2-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl or indol-2-(C<sub>1</sub>-C<sub>4</sub>)alkyl, wherein said preceding R<sub>4</sub> heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, hydroxy or cyano and said substituents are bonded to carbon; or

R<sub>4</sub> is R<sub>15</sub>-carbonyloxymethyl, wherein said R<sub>15</sub> is phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding R<sub>15</sub> rings are optionally mono- or di-substituted independently with halo, amino, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy or trifluoromethyl and said mono- or di-substituents are bonded to carbon;

R<sub>5</sub> is H, methyl, ethyl, n-propyl, hydroxymethyl or hydroxyethyl;

R<sub>6</sub> is carboxy, (C<sub>1</sub>-C<sub>8</sub>)alkoxycarbonyl, benzyloxycarbonyl, C(O)NR<sub>8</sub>R<sub>9</sub> or C(O)R<sub>12</sub>

wherein R<sub>8</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>6</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>6</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkyl, hydroxy or (C<sub>1</sub>-C<sub>8</sub>)alkoxy; and

R<sub>9</sub> is H, cyclo(C<sub>3</sub>-C<sub>8</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>8</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkyl, cyclo(C<sub>4</sub>-C<sub>7</sub>)alkenyl, cyclo(C<sub>3</sub>-C<sub>7</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkoxy, cyclo(C<sub>3</sub>-C<sub>7</sub>)alkyloxy, hydroxy, methylene-perfluorinated (C<sub>1</sub>-C<sub>8</sub>)alkyl, phenyl, or a heterocycle wherein said heterocycle is pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, pyridinyl, piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, benzothiazolyl, benzoxazolyl, benzimidazolyl, thiochromanyl or tetrahydrobenzothiazolyl wherein said heterocycle rings are carbon-nitrogen linked; or

R<sub>9</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>8</sub>)alkoxy wherein said (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>8</sub>)alkoxy is optionally monosubstituted

with cyclo(C<sub>4</sub>-C<sub>7</sub>)alken-1-yl, phenyl, thienyl, pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, piperidinyl, morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, 1,1-dioxothiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl or indolyl and wherein said (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>8</sub>)alkoxy are optionally additionally independently mono- or di-substituted with halo, hydroxy, (C<sub>1</sub>-C<sub>5</sub>)alkoxy, amino, mono-N- or di-N,N- (C<sub>1</sub>-C<sub>5</sub>)alkylamino, cyano, carboxy, or (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl; and

wherein the R<sub>9</sub> rings are optionally mono- or di-substituted independently on carbon with halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, amino(C<sub>1</sub>-C<sub>4</sub>)alkyl, mono-N- or di-N,N- (C<sub>1</sub>-C<sub>4</sub>)alkylamino (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, amino, mono-N- or di-N,N- (C<sub>1</sub>-C<sub>4</sub>)alkylamino, cyano, carboxy, (C<sub>1</sub>-C<sub>5</sub>)alkoxycarbonyl, carbamoyl, formyl or trifluoromethyl and said R<sub>9</sub> rings may optionally be additionally mono- or di-substituted independently with (C<sub>1</sub>-C<sub>5</sub>)alkyl or halo;

with the proviso that no quaternized nitrogen on any R<sub>9</sub> heterocycle is included;

R<sub>12</sub> is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl, azetidin-1-yl, 1,2-oxazinan-2-yl, pyrazolidin-1-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, oxazolidin-3-yl, 3,4-dihydroisoquinolin-2-yl, 1,3-dihydroisoindol-2-yl, 3,4-dihydro-2H-quinol-1-yl, 2,3-dihydro-benzo[1,4]oxazin-4-yl, 2,3-dihydro-benzo[1,4]-thiazine-4-yl, 3,4-dihydro-2H-quinoxalin-1-yl, 3,4-dihydro-benzo[c][1,2]oxazin-1-yl, 1,4-dihydro-

benzo[d][1,2]oxazin-3-yl, 3,4-dihydro-benzo[e][1,2]-  
oxazin-2-yl, 3H-benzo[d]isoxazol-2-yl,  
3H-benzo[c]isoxazol-1-yl or azepan-1-yl,

wherein said  $R_{12}$  rings are optionally mono-, di-  
5 or tri-substituted independently with halo,  $(C_1-C_5)$ alkyl,  
 $(C_1-C_5)$ alkoxy, hydroxy, amino, mono-N- or  
di-N,N- $(C_1-C_5)$ alkylamino, formyl, carboxy, carbamoyl,  
mono-N- or di-N,N- $(C_1-C_5)$ alkylcarbamoyl,  $(C_1-C_6)$ alkoxy  
 $(C_1-C_3)$ alkoxy,  $(C_1-C_5)$ alkoxycarbonyl, benzyloxycarbonyl,  
10  $(C_1-C_5)$ alkoxycarbonyl  $(C_1-C_5)$ alkyl,  
 $(C_1-C_4)$ alkoxycarbonylamino, carboxy  $(C_1-C_5)$ alkyl,  
carbamoyl  $(C_1-C_5)$ alkyl, mono-N- or  
di-N,N- $(C_1-C_5)$ alkylcarbamoyl  $(C_1-C_5)$ alkyl,  
hydroxy  $(C_1-C_5)$ alkyl,  $(C_1-C_4)$ alkoxy  $(C_1-C_4)$ alkyl,  
15 amino  $(C_1-C_4)$ alkyl, mono-N- or  
di-N,N- $(C_1-C_4)$ alkylamino  $(C_1-C_4)$ alkyl, oxo, hydroxyimino or  
 $(C_1-C_6)$ alkoxyimino and wherein no more than two  
substituents are selected from oxo, hydroxyimino or  
 $(C_1-C_6)$ alkoxyimino and oxo, hydroxyimino or  
20  $(C_1-C_6)$ alkoxyimino are on nonaromatic carbon; and  
wherein said  $R_{12}$  rings are optionally  
additionally mono- or di-substituted independently with  
 $(C_1-C_5)$ alkyl or halo;

with the proviso that when  $R_6$  is  
25  $(C_1-C_5)$ alkoxycarbonyl or benzyloxycarbonyl then  $R_1$  is  
5-halo, 5- $(C_1-C_4)$ alkyl or 5-cyano and  $R_4$  is  
(phenyl) (hydroxy)  $(C_1-C_4)$ alkyl,  
(phenyl)  $((C_1-C_4)$ alkoxy)  $(C_1-C_4)$ alkyl, hydroxymethyl or  
Ar  $(C_1-C_2)$ alkyl, wherein Ar is thien-2- or -3-yl, fur-2- or  
30 -3-yl or phenyl wherein said Ar is optionally mono- or  
di-substituted independently with halo; with the provisos  
that when  $R_4$  is benzyl and  $R_5$  is methyl,  $R_{12}$  is not  
4-hydroxy-piperidin-1-yl or when  $R_4$  is benzyl and  $R_5$  is  
methyl  $R_6$  is not  $C(O)N(CH_3)_2$ ;

35 with the proviso that when  $R_1$  and  $R_{10}$  and  $R_{11}$  are  
H,  $R_4$  is not imidazol-4-ylmethyl, 2-phenylethyl or  
2-hydroxy-2-phenylethyl;

with the proviso that when  $R_8$  and  $R_9$  are  $n$ -pentyl,  $R_1$  is 5-chloro, 5-bromo, 5-cyano, 5-( $C_1$ - $C_5$ )alkyl, 5-( $C_1$ - $C_5$ )alkoxy or trifluoromethyl;


with the proviso that when  $R_{12}$  is  
 5 3,4-dihydroisoquinol-2-yl, said 3,4-dihydroisoquinol-2-yl is not substituted with carboxy-( $C_1$ - $C_4$ )alkyl;

with the proviso that when  $R_8$  is H and  $R_9$  is ( $C_1$ - $C_6$ )alkyl,  $R_9$  is not substituted with carboxy or ( $C_1$ - $C_4$ )alkoxycarbonyl on the carbon which is attached to  
 10 the nitrogen atom N of  $NHR_9$ ; and

with the proviso that when  $R_6$  is carboxy and  $R_1$ ,  $R_{10}$ ,  $R_{11}$  and  $R_5$  are all H, then  $R_4$  is not benzyl, H, (phenyl) (hydroxy)methyl, methyl, ethyl or  $n$ -propyl.

Compounds of Formula II are disclosed in  
 15 published Patent Cooperation Treaty Publication number WO 96/39384, the complete disclosure of which is hereby incorporated by reference.

In yet another preferred aspect of the invention, the GPI has the structure of Formula III,  
 20 which is another class of compounds believed to be capable of binding to the indole pocket binding site:



30

Formula III

a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein Formula III has the following substituents:


$R^1$  is ( $C_1$ - $C_4$ )alkyl, ( $C_3$ - $C_7$ )cycloalkyl, phenyl or  
 35 phenyl substituted with up to three ( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy or halogen;

$R^2$  is ( $C_1$ - $C_4$ )alkyl; and

$R^3$  is  $(C_3-C_7)$ cycloalkyl; phenyl; phenyl substituted at the para position with  $(C_1-C_4)$ alkyl, halo, hydroxy $(C_1-C_4)$ alkyl or trifluoromethyl; phenyl substituted at the meta position with fluoro; or phenyl substituted at the ortho position with fluoro.

Compounds of formula III are disclosed more fully in commonly assigned U.S. patent No. 5,998,463, the relevant disclosure of which is incorporated by reference.

In yet another preferred aspect of the invention, the GPI has the structure of Formula IV, which is another class of compounds believed to be capable of binding to the indole pocket binding site:




Formula IV

a stereoisomer, pharmaceutically acceptable salt or prodrug thereof, or a pharmaceutically acceptable salt of the prodrug, wherein Formula IV has the following substituents:


Q is aryl, substituted aryl, heteroaryl, or substituted heteroaryl;  
 each Z and X are independently (C, CH or  $CH_2$ ), N, O or S;  
 $R^1$  is  $NR^a$ ,  $-CH_2-$ , O or S;  
 each - - - is independently a bond or is absent,  
 provided that both - - - are not simultaneously bonds;  
 $R^1$  is hydrogen, halogen,  $-OC_1-C_8$ alkyl,  $-SC_1-C_8$ alkyl,

-C<sub>1</sub>-C<sub>8</sub>alkyl, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHC<sub>1</sub>-C<sub>8</sub>alkyl, -N(C<sub>1</sub>-C<sub>8</sub>alkyl)<sub>2</sub>,  
 -NO<sub>2</sub>, -CN,  
 -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>8</sub>alkyl, -C<sub>2</sub>-C<sub>8</sub>alkenyl, or -C<sub>2</sub>-C<sub>8</sub>alkynyl;  
 each R<sup>a</sup> and R<sup>b</sup> is independently hydrogen or -C<sub>1</sub>-C<sub>8</sub>alkyl;

5



- 10 R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halogen,  
 -C<sub>1</sub>-C<sub>8</sub>alkyl, -CN, -C≡C-Si(CH<sub>3</sub>)<sub>3</sub>,  
 -OC<sub>1</sub>-C<sub>8</sub>alkyl, -SC<sub>1</sub>-C<sub>8</sub>alkyl, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHC<sub>1</sub>-C<sub>8</sub>alkyl,  
 -N(C<sub>1</sub>-C<sub>8</sub>alkyl)<sub>2</sub>,  
 -NO<sub>2</sub>, -CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>8</sub>alkyl, -C<sub>2</sub>-C<sub>8</sub>alkenyl, or  
 15 -C<sub>2</sub>-C<sub>8</sub>alkynyl, or R<sup>2</sup> and R<sup>3</sup> together with the atoms on the  
 ring to which they are attached form a five or six  
 membered ring containing from 0 to 3 heteroatoms and from  
 0 to 2 double bonds;  
 R<sup>4</sup> is -C(=O)-A;  
 20 A is -NR<sup>d</sup>R<sup>d</sup>, -NR<sup>a</sup>CH<sub>2</sub>CH<sub>2</sub>OR<sup>a</sup>,



each R<sup>d</sup> is independently hydrogen, C<sub>1</sub>-C<sub>8</sub>alkyl, C<sub>1</sub>-C<sub>8</sub>alkoxy,  
 aryl, substituted aryl, heteroaryl, or substituted  
 heteroaryl;

- 30 each R<sup>c</sup> is independently hydrogen, -C(=O)OR<sup>a</sup>, -OR<sup>a</sup>, -SR<sup>a</sup>,  
 or -NR<sup>a</sup>R<sup>a</sup>; and  
 each n is independently 1-3.

35 Compounds of Formula IV are disclosed in commonly  
 assigned U.S. Provisional Patent Application Serial  
 No. 60/157,148, filed September 30, 1999, the relevant  
 disclosure of which is incorporated by reference.



In an especially preferred embodiment, the GPI is selected from one of the following compounds of Formula I:

5 5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-dimethylcarbamoylmethyl)-2-phenyl-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-methoxy-methylcarbamoylmethyl)-2-phenyl-ethyl]-amide;

10

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3S)-hydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide;

15

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide;

20

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4R)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide; and

25

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-morpholin-4-yl-3-oxo-propyl]-amide.

In another especially preferred embodiment, the GPI is selected from one of the following compounds of Formula II:

30

5-chloro-1H-indole-2-carboxylic acid [2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

35

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide;

5 5-chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluorobenzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide;

5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide;

10 5-chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide; and

5-chloro-1H-indole-2-carboxylic acid [2-(1-oxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide.

In another especially preferred embodiment, the GPI is selected from one of the following compounds of Formula III:

20 5-acetyl-1-ethyl-2,3-dihydro-2-oxo-N-[3-[(phenylamino)carbonyl]phenyl]-1H-Indole-3-carboxamide;

5-acetyl-N-[3-[(cyclohexylamino)carbonyl]phenyl]-1-ethyl-2,3-dihydro-2-oxo-1H-Indole-3-carboxamide; and

5-acetyl-N-[3-[[4-bromophenyl]amino]carbonyl]phenyl]-2,3-dihydro-1-methyl-2-oxo-1H-Indole-3-carboxamide.

30 In another especially preferred embodiment, the GPI is selected from one of the following compounds of Formula IV:

2-Chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide; and

2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide.

5

## CONCENTRATION-ENHANCING POLYMERS

Concentration-enhancing polymers suitable for use in the compositions of the present invention should be inert, in the sense that they do not chemically react with the GPI in an adverse manner, are pharmaceutically acceptable, and have at least some solubility in aqueous solution at physiologically relevant pHs (e.g. 1-8). The polymer can be neutral or ionizable, and should have an aqueous-solubility of at least 0.1 mg/mL over at least a portion of the pH range of 1-8. The polymer is a "concentration-enhancing polymer," meaning that it meets at least one, and more preferably both, of the following conditions. The first condition is that the concentration-enhancing polymer increases the MDC of the GPI in the environment of use relative to a control composition consisting of an equivalent amount of the GPI but no polymer. That is, once the composition is introduced into an environment of use, the polymer increases the aqueous concentration of GPI relative to the control composition. Preferably, the polymer increases the MDC of the GPI in aqueous solution by at least 1.25-fold relative to a control composition, and more preferably by at least 2-fold and most preferably by at least 3-fold. The second condition is that the concentration-enhancing polymer increases the AUC of the GPI in the environment of use relative to a control composition consisting of GPI but no polymer as described above. That is, in the environment of use, the composition comprising the GPI and the concentration-enhancing polymer provides an area under the concentration versus time curve (AUC) for any period of 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction

to the use environment that is at least 1.25-fold that of a control composition comprising an equivalent quantity of GPI but no polymer.

Concentration-enhancing polymers suitable for use with the present invention may be cellulosic or non-cellulosic. The polymers may be neutral or ionizable in aqueous solution. Of these, ionizable and cellulosic polymers are preferred, with ionizable cellulosic polymers being more preferred.

A preferred class of polymers comprises polymers that are "amphiphilic" in nature, meaning that the polymer has hydrophobic and hydrophilic portions. Hydrophobic groups may comprise groups such as aliphatic or aromatic hydrocarbon groups. Hydrophilic groups may comprise either ionizable or non-ionizable groups that are capable of hydrogen bonding such as hydroxyls, carboxylic acids, esters, amines or amides.

Amphiphilic and/or ionizable polymers are preferred because it is believed that such polymers may tend to have relatively strong interactions with the GPI and may promote the formation of the various types of polymer/drug assemblies in the use environment as described previously. In addition, the repulsion of the like charges of the ionized groups of such polymers may serve to limit the size of the polymer/drug assemblies to the nanometer or submicron scale. For example, while not wishing to be bound by a particular theory, such polymer/drug assemblies may comprise hydrophobic GPI clusters surrounded by the polymer with the polymer's hydrophobic regions turned inward towards the GPI and the hydrophilic regions of the polymer turned outward toward the aqueous environment. Alternatively, depending on the specific chemical nature of the GPI, the ionized functional groups of the polymer may associate, for example, via ion pairing or hydrogen bonds, with ionic or polar groups of the GPI. In the case of ionizable polymers, the hydrophilic regions of the polymer would

include the ionized functional groups. Such polymer/drug assemblies in solution may well resemble charged polymeric micellar-like structures. In any case, regardless of the mechanism of action, the inventors have  
5 observed that such amphiphilic polymers, particularly ionizable cellulosic polymers, have been shown to improve the MDC and/or AUC of GPI in aqueous solution relative to control compositions free from such polymers.

Surprisingly, such amphiphilic polymers can  
10 greatly enhance the maximum concentration of GPI obtained when an amorphous form of the GPI is dosed to a use environment. In addition, such amphiphilic polymers interact with the GPI to prevent the precipitation or crystallization of the GPI from solution despite its  
15 concentration being substantially above its equilibrium concentration. In particular, when the preferred compositions are solid amorphous dispersions of the GPI and the concentration-enhancing polymer, the compositions provide a greatly enhanced drug concentration,  
20 particularly when the dispersions are substantially homogeneous. The maximum drug concentration may be 2-fold and often up to 10-fold the equilibrium concentration of the crystalline GPI. Such enhanced GPI concentrations in turn lead to substantially enhanced  
25 relative bioavailability for the GPI.

One class of polymers suitable for use with the present invention comprises neutral non-cellulosic polymers. Exemplary polymers include: vinyl polymers and copolymers having substituents of hydroxyl,  
30 alkylacyloxy, and cyclicamido; polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed (vinyl acetate) form; polyvinyl alcohol polyvinyl acetate copolymers; polyvinyl pyrrolidone; and polyethylene polyvinyl alcohol copolymers.

35 Another class of polymers suitable for use with the present invention comprises ionizable non-cellulosic polymers. Exemplary polymers include: carboxylic acid-

functionalized vinyl polymers, such as the carboxylic acid functionalized polymethacrylates and carboxylic acid functionalized polyacrylates such as the EUDRAGITS® manufactured by Rohm Tech Inc., of Malden, Massachusetts; amine-functionalized polyacrylates and polymethacrylates; proteins; and carboxylic acid functionalized starches such as starch glycolate.

Non-cellulosic polymers that are amphiphilic are copolymers of a relatively hydrophilic and a relatively hydrophobic monomer. Examples include acrylate and methacrylate copolymers. Exemplary commercial grades of such copolymers include the EUDRAGITS, which are copolymers of methacrylates and acrylates.

A preferred class of polymers comprises ionizable and neutral cellulosic polymers with at least one ester- and/or ether- linked substituent in which the polymer has a degree of substitution of at least 0.1 for each substituent. It should be noted that in the polymer nomenclature used herein, ether-linked substituents are recited prior to "cellulose" as the moiety attached to the ether group; for example, "ethylbenzoic acid cellulose" has ethoxybenzoic acid substituents. Analogously, ester-linked substituents are recited after "cellulose" as the carboxylate; for example, "cellulose phthalate" has one carboxylic acid of each phthalate moiety ester-linked to the polymer and the other carboxylic acid unreacted.

It should also be noted that a polymer name such as "cellulose acetate phthalate" (CAP) refers to any of the family of cellulosic polymers that have acetate and phthalate groups attached via ester linkages to a significant fraction of the cellulosic polymer's hydroxyl groups. Generally, the degree of substitution of each substituent group can range from 0.1 to 2.9 as long as the other criteria of the polymer are met. "Degree of substitution" refers to the average number of the three

hydroxyls per saccharide repeat unit on the cellulose chain that have been substituted. For example, if all of the hydroxyls on the cellulose chain have been phthalate substituted, the phthalate degree of substitution is 3.

5 Also included within each polymer family type are cellulosic polymers that have additional substituents added in relatively small amounts that do not substantially alter the performance of the polymer.

Amphiphilic cellulosics may be prepared by  
10 substituting the cellulosic at any or all of the 3 hydroxyl substituents present on each saccharide repeat unit with at least one relatively hydrophobic substituent. Hydrophobic substituents may be essentially any substituent that, if substituted to a high enough  
15 level or degree of substitution, can render the cellulosic polymer essentially aqueous insoluble. Hydrophilic regions of the polymer can be either those portions that are relatively unsubstituted, since the unsubstituted hydroxyls are themselves relatively  
20 hydrophilic, or those regions that are substituted with hydrophilic substituents. Examples of hydrophobic substituents include ether-linked alkyl groups such as methyl, ethyl, propyl, butyl, etc.; or ester-linked alkyl groups such as acetate, propionate, butyrate, etc.; and  
25 ether- and/or ester-linked aryl groups such as phenyl, benzoate, or phenylate. Hydrophilic groups include ether- or ester-linked nonionizable groups such as the hydroxy alkyl substituents hydroxyethyl, hydroxypropyl, and the alkyl ether groups such as ethoxyethoxy or  
30 methoxyethoxy. Particularly preferred hydrophilic substituents are those that are ether- or ester-linked ionizable groups such as carboxylic acids, thiocarboxylic acids, substituted phenoxy groups, amines, phosphates or sulfonates.

35 One class of cellulosic polymers comprises neutral polymers, meaning that the polymers are substantially non-ionizable in aqueous solution. Such

polymers contain non-ionizable substituents, which may be either ether-linked or ester-linked. Exemplary ether-linked non-ionizable substituents include: alkyl groups, such as methyl, ethyl, propyl, butyl, etc.; hydroxy alkyl groups such as hydroxymethyl, hydroxyethyl, hydroxypropyl, etc.; and aryl groups such as phenyl. Exemplary ester-linked non-ionizable groups include: alkyl groups, such as acetate, propionate, butyrate, etc.; and aryl groups such as phenylate. However, when aryl groups are included, the polymer may need to include a sufficient amount of a hydrophilic substituent so that the polymer has at least some water solubility at any physiologically relevant pH of from 1 to 8.

Exemplary non-ionizable polymers that may be used as the polymer include: hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose.

A preferred set of neutral cellulosic polymers are those that are amphiphilic. Exemplary polymers include hydroxypropyl methyl cellulose and hydroxypropyl cellulose acetate, where cellulosic repeat units that have relatively high numbers of methyl or acetate substituents relative to the unsubstituted hydroxyl or hydroxypropyl substituents constitute hydrophobic regions relative to other repeat units on the polymer.

A preferred class of cellulosic polymers comprises polymers that are at least partially ionizable at physiologically relevant pH and include at least one ionizable substituent, which may be either ether-linked or ester-linked. Exemplary ether-linked ionizable substituents include: carboxylic acids, such as acetic acid, propionic acid, benzoic acid, salicylic acid, alkoxybenzoic acids such as ethoxybenzoic acid or propoxybenzoic acid, the various isomers of alkoxyphthalic acid such as ethoxyphthalic acid and



ethoxyisophthalic acid, the various isomers of  
alkoxynicotinic acid such as ethoxynicotinic acid, and  
the various isomers of picolinic acid such as  
ethoxypicolinic acid, etc.; thiocarboxylic acids, such as  
5 thioacetic acid; substituted phenoxy groups, such as  
hydroxyphenoxy, etc.; amines, such as aminoethoxy,  
diethylaminoethoxy, trimethylaminoethoxy, etc.;  
phosphates, such as phosphate ethoxy; and sulfonates,  
such as sulphonate ethoxy. Exemplary ester linked  
10 ionizable substituents include: carboxylic acids, such  
as succinate, citrate, phthalate, terephthalate,  
isophthalate, trimellitate, and the various isomers of  
pyridinedicarboxylic acid, etc.; thiocarboxylic acids,  
such as thiosuccinate; substituted phenoxy groups, such  
15 as amino salicylic acid; amines, such as natural or  
synthetic amino acids, such as alanine or phenylalanine;  
phosphates, such as acetyl phosphate; and sulfonates,  
such as acetyl sulfonate. For aromatic-substituted  
polymers to also have the requisite aqueous solubility,  
20 it is also desirable that sufficient hydrophilic groups  
such as hydroxypropyl or carboxylic acid functional  
groups be attached to the polymer to render the polymer  
aqueous soluble at least at pH values where any ionizable  
groups are ionized. In some cases, the aromatic group  
25 may itself be ionizable, such as phthalate or  
trimellitate substituents.

Exemplary ionizable cellulosic polymers that  
are at least partially ionized at physiologically  
relevant pHs include: hydroxypropyl methyl cellulose  
30 acetate succinate, hydroxypropyl methyl cellulose  
succinate, hydroxypropyl cellulose acetate succinate,  
hydroxyethyl methyl cellulose succinate, hydroxyethyl  
cellulose acetate succinate, hydroxypropyl methyl  
cellulose phthalate, hydroxyethyl methyl cellulose  
35 acetate succinate, hydroxyethyl methyl cellulose acetate  
phthalate, carboxyethyl cellulose, carboxymethyl  
cellulose, cellulose acetate phthalate, methyl cellulose

acetate phthalate, ethyl cellulose acetate phthalate,  
hydroxypropyl cellulose acetate phthalate, hydroxypropyl  
methyl cellulose acetate phthalate, hydroxypropyl  
cellulose acetate phthalate succinate, hydroxypropyl  
5 methyl cellulose acetate succinate phthalate,  
hydroxypropyl methyl cellulose succinate phthalate,  
cellulose propionate phthalate, hydroxypropyl cellulose  
butyrate phthalate, cellulose acetate trimellitate,  
methyl cellulose acetate trimellitate, ethyl cellulose  
10 acetate trimellitate, hydroxypropyl cellulose acetate  
trimellitate, hydroxypropyl methyl cellulose acetate  
trimellitate, hydroxypropyl cellulose acetate  
trimellitate succinate, cellulose propionate  
trimellitate, cellulose butyrate trimellitate, cellulose  
15 acetate terephthalate, cellulose acetate isophthalate,  
cellulose acetate pyridinedicarboxylate, salicylic acid  
cellulose acetate, hydroxypropyl salicylic acid cellulose  
acetate, ethylbenzoic acid cellulose acetate,  
hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl  
20 phthalic acid cellulose acetate, ethyl nicotinic acid  
cellulose acetate, and ethyl picolinic acid cellulose  
acetate.

Exemplary cellulosic polymers that meet the  
definition of amphiphilic, having hydrophilic and  
25 hydrophobic regions include polymers such as cellulose  
acetate phthalate and cellulose acetate trimellitate  
where the cellulosic repeat units that have one or more  
acetate substituents are hydrophobic relative to those  
that have no acetate substituents or have one or more  
30 ionized phthalate or trimellitate substituents.

A particularly desirable subset of cellulosic  
ionizable polymers are those that possess both a  
carboxylic acid functional aromatic substituent and an  
alkylate substituent and thus are amphiphilic. Exemplary  
35 polymers include cellulose acetate phthalate, methyl  
cellulose acetate phthalate, ethyl cellulose acetate  
phthalate, hydroxypropyl cellulose acetate phthalate,

hydroxypropyl methyl cellulose phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate.

Another particularly desirable subset of cellulosic ionizable polymers are those that possess a non-aromatic carboxylate substituent. Exemplary polymers include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, and hydroxyethyl cellulose acetate succinate.

Especially preferred polymers are hydroxypropyl methyl cellulose acetate succinate (HPMCAS), hydroxypropyl methyl cellulose phthalate (HPMCP), cellulose acetate phthalate (CAP), cellulose acetate trimellitate (CAT), methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate and cellulose acetate isophthalate. The most preferred polymers are hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl

cellulose phthalate, cellulose acetate phthalate, and cellulose acetate trimellitate.

While specific polymers have been discussed as being suitable for use in the mixtures of the present invention, blends of such polymers may also be suitable. Thus the term "polymer" is intended to include blends of polymers in addition to a single species of polymer.

To obtain the best performance, particularly upon storage for long times prior to use, it is preferred that the GPI remain, to the extent possible, in the amorphous state. The inventors have found that this is best achieved when the glass-transition temperature,  $T_g$ , of the amorphous GPI material is substantially above the storage temperature of the composition. In particular, it is preferable that the  $T_g$  of the amorphous state of the GPI be at least 40°C and preferably greater than 60°C. For those aspects of the invention in which the composition is a solid, substantially amorphous dispersion of GPI in the concentration-enhancing polymer and in which the GPI itself has a relatively low  $T_g$  (about 70°C or less) it is preferred that the concentration-enhancing polymer have a  $T_g$  of at least 40°C, preferably at least 70°C and more preferably greater than 100°C. Exemplary high  $T_g$  polymers include HPMCAS, HPMCP, CAP, CAT, and other cellulosics that have alkylate or aromatic substituents or both alkylate and aromatic substituents. In addition, the preferred polymers listed above, that is amphiphilic cellulosic polymers, tend to have greater concentration-enhancing properties relative to the other polymers of the present invention. For any particular GPI, the amphiphilic cellulosic with the best concentration-enhancing properties may vary. However, the inventors have found that generally those that have ionizable substituents tend to perform best. In vitro tests of compositions with such polymers tend to have higher MDC and AUC values than compositions with other polymers of the invention. Often such compositions have

MDC and AUC values that are more than 4-fold and in some cases more than 8-fold that of a control composition.

#### PREPARATION OF COMPOSITIONS

5                   Compositions may comprise a physical mixture of GPI and concentration-enhancing polymer or a dispersion of GPI and polymer. Preferably, the compositions are formed such that at least a major portion (at least 60%) of the GPI is in the amorphous state. In cases where the  
10 composition is a physical mixture of amorphous GPI and polymer the amorphous GPI may be made by any known process. Generally the amorphous form of the GPI is made by (1) melting the drug followed by rapid cooling (e.g., melt-congeal process); (2) dissolution of the drug in a  
15 solvent followed by precipitation or evaporation (e.g., spray drying, spray coating); or (3) mechanical processing of the drug (e.g., extrusion, ball milling). Various combinations of heat (as in melt processes), solvent and mechanical force may be used to generate the  
20 amorphous GPI.

                  Dispersions of the GPI and concentration-enhancing polymer may be made according to any known process which results in at least a major portion (at least 60%) of the GPI being in the amorphous state.  
25 Exemplary mechanical processes include milling and extrusion; melt processes include high temperature fusion, solvent modified fusion and melt-congeal processes; and solvent processes include non-solvent precipitation, spray coating and spray-drying. Although  
30 the dispersions of the present invention may be made by any of these processes, the dispersions generally have their maximum bioavailability and stability when the GPI is dispersed in the polymer such that it is substantially amorphous and substantially homogeneously distributed  
35 throughout the polymer. Although in some cases such substantially amorphous and substantially homogeneous dispersions may be made by any of these methods, it has

been found that such dispersions are preferably formed by "solvent processing," which consists of dissolution of the GPI and one or more polymers in a common solvent. "Common" here means that the solvent, which can be a mixture of compounds, will simultaneously dissolve the drug and the polymer(s). After both the GPI and the polymer have been dissolved, the solvent is rapidly removed by evaporation or by mixing with a non-solvent. Exemplary processes are spray-drying, spray-coating (pan-coating, fluidized bed coating, etc.), and precipitation by rapid mixing of the polymer and drug solution with CO<sub>2</sub>, water, or some other non-solvent. Preferably, removal of the solvent results in a solid dispersion which is substantially homogeneous. As described previously, in such substantially homogeneous dispersions, the GPI is dispersed as homogeneously as possible throughout the polymer and can be thought of as a solid solution of GPI dispersed in the polymer(s). When the resulting dispersion constitutes a solid solution of GPI in polymer, the dispersion may be thermodynamically stable, meaning that the concentration of GPI in the polymer is at or below its equilibrium value, or it may be considered a supersaturated solid solution where the GPI concentration in the dispersion polymer(s) is above its equilibrium value.

The solvent may be removed through the process of spray-drying. The term spray-drying is used conventionally and broadly refers to processes involving breaking up liquid mixtures into small droplets (atomization) and rapidly removing solvent from the mixture in a container (spray-drying apparatus) where there is a strong driving force for evaporation of solvent from the droplets. The strong driving force for solvent evaporation is generally provided by maintaining the partial pressure of solvent in the spray-drying apparatus well below the vapor pressure of the solvent at the temperature of the drying droplets. This is

accomplished by either (1) maintaining the pressure in the spray-drying apparatus at a partial vacuum (e.g., 0.01 to 0.50 atm); (2) mixing the liquid droplets with a warm drying gas; or (3) both. In addition, at least a portion of the heat required for evaporation of solvent may be provided by heating the spray solution.

Solvents suitable for spray-drying can be any organic compound in which the GPI and polymer are mutually soluble. Preferably, the solvent is also volatile with a boiling point of 150°C or less. In addition, the solvent should have relatively low toxicity and be removed from the dispersion to a level that is acceptable according to The International Committee on Harmonization (ICH) guidelines. Removal of solvent to this level may require a processing step such as tray-drying subsequent to the spray-drying or spray-coating process. Preferred solvents include alcohols such as methanol, ethanol, n-propanol, iso-propanol, and butanol; ketones such as acetone, methyl ethyl ketone and methyl iso-butyl ketone; esters such as ethyl acetate and propylacetate; and various other solvents such as acetonitrile, methylene chloride, toluene, and 1,1,1-trichloroethane. Lower volatility solvents such as dimethyl acetamide or dimethylsulfoxide can also be used. Mixtures of solvents, such as 50% methanol and 50% acetone, can also be used, as can mixtures with water as long as the polymer and GPI are sufficiently soluble to make the spray-drying process practicable. Generally, non-aqueous solvents are preferred meaning that the solvent comprises less than about 40 wt% water. However, for certain GPIs, it has been found that addition of a small amount of water, typically about 5 wt% to about 35 wt%, to a solvent such as acetone may actually increase the solubility of the GPI in the solvent; relative to that in the absence of water. In such cases, or to enhance the polymer solubility, addition of water may even be preferred.

Generally, the temperature and flow rate of the drying gas is chosen so that the polymer/drug-solution droplets are dry enough by the time they reach the wall of the apparatus that they are essentially solid, and so that they form a fine powder and do not stick to the apparatus wall. The actual length of time to achieve this level of dryness depends on the size of the droplets. Droplet sizes generally range from 1  $\mu\text{m}$  to 500  $\mu\text{m}$  in diameter, with 5 to 100  $\mu\text{m}$  being more typical. The large surface-to-volume ratio of the droplets and the large driving force for evaporation of solvent leads to actual drying times of a few seconds or less, and more typically less than 0.1 second. This rapid drying is often critical to the particles maintaining a uniform, homogeneous dispersion instead of separating into drug-rich and polymer-rich phases. Solidification times should be less than 100 seconds, preferably less than a few seconds, and more preferably less than 1 second. In general, to achieve this rapid solidification of the GPI/polymer solution, it is preferred that the size of droplets formed during the spray-drying process are less than 100  $\mu\text{m}$  in diameter, preferably less than 50  $\mu\text{m}$  in diameter, and more preferably less than 25  $\mu\text{m}$  in diameter. The resultant solid particles thus formed are generally less than 100  $\mu\text{m}$  in diameter, and preferably less than 50  $\mu\text{m}$  in diameter, and more preferably less than 25  $\mu\text{m}$  in diameter. Typically, particles are 1 to 20  $\mu\text{m}$  in diameter.

Following solidification, the solid powder typically stays in the spray-drying chamber for about 5 to 60 seconds, further evaporating solvent from the solid powder. The final solvent content of the solid dispersion as it exits the dryer should be low, since this reduces the mobility of GPI molecules in the dispersion, thereby improving its stability. Generally, the solvent content of the dispersion as it leaves the spray-drying chamber should be less than 10 wt% and



preferably less than 2 wt%. In some cases, it may be preferable to spray a solvent or a solution of a polymer or other excipient into the spray-drying chamber to form granules, so long as the dispersion is not adversely affected.

Spray-drying processes and spray-drying equipment are described generally in Perry's *Chemical Engineers' Handbook*, Sixth Edition (R. H. Perry, D. W. Green, J. O. Maloney, eds.) McGraw-Hill Book Co. 1984, pages 20-54 to 20-57. More details on spray-drying processes and equipment are reviewed by Marshall "Atomization and Spray-Drying," 50 *Chem. Eng. Prog. Monogr. Series 2* (1954).

Where the composition is a simple physical mixture, the composition may be prepared by dry- or wet-mixing the drug or drug mixture with the polymer to form the composition. Mixing processes include physical processing as well as wet-granulation and coating processes. Any conventional mixing method may be used, including those that substantially convert the drug and polymer to a molecular dispersion.

For example, mixing methods include convective mixing, shear mixing, or diffusive mixing. Convective mixing involves moving a relatively large mass of material from one part of a powder bed to another, by means of blades or paddles, revolving screw, or an inversion of the powder bed. Shear mixing occurs when slip planes are formed in the material to be mixed. Diffusive mixing involves an exchange of position by single particles. These mixing processes can be performed using equipment in batch or continuous mode. Tumbling mixers (e.g., twin-shell) are commonly used equipment for batch processing. Continuous mixing can be used to improve composition uniformity.

Milling may also be employed to prepare the compositions of the present invention. Milling is the

mechanical process of reducing the particle size of solids (comminution). The most common types of milling equipment are the rotary cutter, the hammer, the roller and fluid energy mills. Equipment choice depends on the characteristics of the ingredients in the drug form (e.g., soft, abrasive, or friable). Wet- or dry-milling techniques can be chosen for several of these processes, also depending on the characteristics of the ingredients (e.g., drug stability in solvent). The milling process may serve simultaneously as a mixing process if the feed materials are heterogeneous. Conventional mixing and milling processes suitable for use in the present invention are discussed more fully in Lachman, et al., *The Theory and Practice of Industrial Pharmacy* (3d Ed. 1986). The components of the compositions of this invention may also be combined by dry- or wet-granulating processes.

In addition to the physical mixtures described above, the compositions of the present invention may constitute any device or collection of devices that accomplishes the objective of delivering to the use environment both the GPI and the concentration-enhancing polymer. Thus, in the case of oral administration to an animal, the dosage form may constitute a layered tablet wherein one or more layers comprise the amorphous GPI and one or more other layers comprise the polymer. Alternatively, the dosage form may be a coated tablet wherein the tablet core comprises the GPI and the coating comprises the concentration-enhancing polymer. In addition, the GPI and the polymer may even be present in different dosage forms such as tablets or beads and may be administered simultaneously or separately as long as both the GPI and polymer are administered in such a way that the GPI and polymer can come into contact in the use environment. When the GPI and the polymer are administered separately it is generally preferable to deliver the polymer prior to the GPI.

The amount of concentration-enhancing polymer relative to the amount of GPI present in the mixtures of the present invention depends on the GPI and polymer and may vary widely from a GPI-to-polymer weight ratio of  
5 from 0.01 to about 4 (e.g., 1 wt% GPI to 80 wt% GPI).

However, in most cases it is preferred that the GPI-to-polymer ratio is greater than about 0.05 (4.8 wt% GPI) and less than about 2.5 (71 wt% GPI). Often the enhancement in GPI concentration or relative  
10 bioavailability that is observed increases as the GPI-to-polymer ratio decreases from a value of about 1 (50 wt% GPI) to a value of about 0.11 (10 wt% GPI). The maximum GPI:polymer ratio that yields satisfactory results varies from GPI to GPI and is best determined in *in vitro*  
15 dissolution tests and/or *in vivo* bioavailability tests.

It should be noted that this level of concentration-enhancing polymer is usually substantially greater and often much greater than the amount of polymer conventionally included in dosage forms for other uses  
20 such as binders or coatings. Thus, it is preferred in the compositions of this invention that there be included sufficient concentration-enhancing polymer that the compositions meet the *in vitro* MDC and AUC criteria and *in vivo* bioavailability criterion previously set forth.

25 In general, to maximize the GPI concentration or relative bioavailability of the GPI, lower GPI-to-polymer ratios are preferred. At low GPI-to-polymer ratios, there is sufficient polymer available in solution to ensure the inhibition of the precipitation or  
30 crystallization of GPI from solution and, thus, the average concentration of GPI is much higher. For high GPI-to-polymer ratios, not enough polymer may be present in solution and GPI precipitation or crystallization of the GPI may occur more readily. In addition, the amount  
35 of concentration-enhancing polymer that can be used in a dosage form is often limited by the total mass

requirements of the dosage form. For example, when oral dosing to a human is desired, at low GPI-to-polymer ratios the total mass of drug and polymer may be unacceptably large for delivery of the desired dose in a single tablet or capsule. Thus, it is often necessary to use GPI-to-polymer ratios that are less than optimum in specific dosage forms to provide a sufficient GPI dose in a dosage form that is small enough to be easily delivered to a use environment.

10

#### EXCIPIENTS AND DOSAGE FORMS

Although the key ingredients present in the compositions of the present invention are simply the GPI to be delivered and the concentration-enhancing polymer(s), the inclusion of other excipients in the composition may be useful. These excipients may be utilized with the GPI/polymer mixture in order to formulate the mixture into tablets, capsules, suspensions, powders for suspension, creams, transdermal patches, depots, and the like. The amorphous GPI and polymer can be added to other dosage form ingredients in essentially any manner that does not substantially alter the GPI. In addition, as described above, the GPI and the polymer may be mixed with excipients separately to form different beads, or layers, or coatings, or cores or even separate dosage forms.

One very useful class of excipients is surfactants. Suitable surfactants include fatty acid and alkyl sulfonates; commercial surfactants such as benzalkonium chloride (HYAMINE® 1622, available from Lonza, Inc., Fairlawn, New Jersey); dioctyl sodium sulfosuccinate, DOCUSATE SODIUM™ (available from Mallinckrodt Spec. Chem., St. Louis, Missouri); polyoxyethylene sorbitan fatty acid esters (TWEEN®, available from ICI Americas Inc., Wilmington, Delaware; LIPOSORB® P-20 available from Lipochem Inc., Patterson New Jersey; CAPMUL® POE-0 available from Abitec Corp.,

35

Janesville, Wisconsin), and natural surfactants such as sodium taurocholic acid, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine, lecithin, and other phospholipids and mono- and diglycerides. Such materials can  
5 advantageously be employed to increase the rate of dissolution by facilitating wetting, thereby increasing the maximum dissolved concentration, and also to inhibit crystallization or precipitation of drug by interacting with the dissolved drug by mechanisms such as  
10 complexation, formation of inclusion complexes, formation of micelles or adsorbing to the surface of solid drug, crystalline or amorphous. These surfactants may comprise up to 5 wt% of the composition.

The addition of pH modifiers such as acids, bases, or buffers may also be beneficial, retarding the  
15 dissolution of the composition (e.g., acids such as citric acid or succinic acid when the concentration-enhancing polymer is anionic) or, alternatively, enhancing the rate of dissolution of the composition (e.g., bases such as sodium acetate or amines when the  
20 polymer is anionic).

Conventional matrix materials, complexing agents, solubilizers, fillers, disintegrating agents (disintegrants), or binders may also be added as part of  
25 the composition itself or added by granulation via wet or mechanical or other means. These materials may comprise up to 90 wt% of the composition.

Examples of matrix materials, fillers, or diluents include lactose, mannitol, xylitol,  
30 microcrystalline cellulose, calcium diphosphate, and starch.

Examples of disintegrants include sodium starch glycolate, sodium alginate, carboxy methyl cellulose sodium, methyl cellulose, and croscarmellose sodium.

35 Examples of binders include methyl cellulose, microcrystalline cellulose, starch, and gums such as guar gum, and tragacanth.

Examples of lubricants include magnesium stearate and calcium stearate.

Other conventional excipients may be employed in the compositions of this invention, including those  
5 excipients well-known in the art. Generally, excipients such as pigments, lubricants, flavorants, and so forth may be used for customary purposes and in typical amounts without adversely affecting the properties of the  
10 compositions. These excipients may be utilized in order to formulate the composition into tablets, capsules, suspensions, powders for suspension, creams, transdermal patches, and the like.

Compositions of this invention may be used in a wide variety of dosage forms for administration of GPIs.  
15 Exemplary dosage forms are powders or granules that may be taken orally either dry or reconstituted by addition of water or other liquids to form a paste, slurry, suspension or solution; tablets; capsules; multiparticulates; and pills. Various additives may be  
20 mixed, ground, or granulated with the compositions of this invention to form a material suitable for the above dosage forms.

The compositions of the present invention may be formulated in various forms such that they are  
25 delivered as a suspension of particles in a liquid vehicle. Such suspensions may be formulated as a liquid or paste at the time of manufacture, or they may be formulated as a dry powder with a liquid, typically water, added at a later time but prior to oral  
30 administration. Such powders that are constituted into a suspension are often termed sachets or oral powder for constitution (OPC) formulations. Such dosage forms can be formulated and reconstituted via any known procedure. The simplest approach is to formulate the dosage form as  
35 a dry powder that is reconstituted by simply adding water and agitating. Alternatively, the dosage form may be formulated as a liquid and a dry powder that are combined

and agitated to form the oral suspension. In yet another embodiment, the dosage form can be formulated as two powders which are reconstituted by first adding water to one powder to form a solution to which the second powder is combined with agitation to form the suspension.

Generally, it is preferred that the dispersion of GPI or amorphous form of GPI be formulated for long-term storage in the dry state as this promotes the chemical and physical stability of the GPI. Various excipients and additives are combined with the compositions of the present invention to form the dosage form. For example, it may be desirable to add some or all of the following: preservatives such as sulfites (an antioxidant), benzalkonium chloride, methyl paraben, propyl paraben, benzyl alcohol or sodium benzoate; suspending agents or thickeners such as xanthan gum, starch, guar gum, sodium alginate, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, hydroxypropyl methyl cellulose, polyacrylic acid, silica gel, aluminum silicate, magnesium silicate, or titanium dioxide; anticaking agents or fillers such as silicon oxide, or lactose; flavorants such as natural or artificial flavors; sweeteners such as sugars such as sucrose, lactose, or sorbitol as well as artificial sweeteners such as aspartame or saccharin; wetting agents or surfactants such as various grades of polysorbate, docusate sodium, or sodium lauryl sulfate; solubilizers such as ethanol propylene glycol or polyethylene glycol; coloring agents such as FD and C Red No. 3 or FD and C Blue No. 1; and pH modifiers or buffers such as carboxylic acids (including citric acid, ascorbic acid, lactic acid, and succinic acid), various salts of carboxylic acids, amino acids such as glycine or alanine, various phosphate, sulfate and carbonate salts such as trisodium phosphate, sodium bicarbonate or potassium bisulfate, and bases such as amino glucose or triethanol amine.

A preferred additive to such formulations is additional concentration-enhancing polymer which may act as a thickener or suspending agent as well as to enhance the concentration of GPI in the environment of use and may also act to prevent or retard precipitation or crystallization of GPI from solution. Such preferred additives are hydroxyethyl cellulose, hydroxypropyl cellulose, and hydroxypropyl methyl cellulose. In particular, the salts of carboxylic acid functional polymers such as cellulose acetate phthalate, hydroxypropyl methyl cellulose acetate succinate, and carboxymethyl cellulose are useful in this regard. Such polymers may be added in their salt forms or the salt form may be formed *in situ* during reconstitution by adding a base such as trisodium phosphate and the acid form of such polymers.

In some cases, the overall dosage form or particles, granules or beads that make up the dosage form may have superior performance if coated with an enteric polymer to prevent or retard dissolution until the dosage form leaves the stomach. Exemplary enteric coating materials include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, carboxylic acid-functionalized polymethacrylates, and carboxylic acid-functionalized polyacrylate.

Compositions of this invention may be administered in a controlled release dosage form. In one such dosage form, the composition of the GPI and polymer is incorporated into an erodible polymeric matrix device. By an erodible matrix is meant aqueous-erodible or water-swallowable or aqueous-soluble in the sense of being either erodible or swellable or dissolvable in pure water or requiring the presence of an acid or base to ionize the polymeric matrix sufficiently to cause erosion or dissolution. When contacted with the aqueous environment



of use, the erodible polymeric matrix imbibes water and forms an aqueous-swollen gel or "matrix" that entraps the mixture of GPI and polymer. The aqueous-swollen matrix gradually erodes, swells, disintegrates or dissolves in the environment of use, thereby controlling the release of the drug mixture to the environment of use. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,059 filed January 31, 2000 which claimed the benefit of priority of provisional patent application Serial No. 60/119,400 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions of the present invention may be administered by or incorporated into a non-erodible matrix device.

Alternatively, the drug mixture of the invention may be delivered using a coated osmotic controlled release dosage form. This dosage form has two components: (a) the core which contains an osmotic agent and the GPI and the concentration-enhancing polymer; and (b) a non-dissolving and non-eroding coating surrounding the core, the coating controlling the influx of water to the core from an aqueous environment of use so as to cause drug release by extrusion of some or all of the core to the environment of use. The GPI and the concentration-enhancing polymer may be homogeneously distributed throughout the core or they may be partially or completely segregated in separate regions of the core. The osmotic agent contained in the core of this device may be an aqueous-swellaable hydrophilic polymer, osmogen, or osmagent. The coating is preferably polymeric, aqueous-permeable, and has at least one delivery port. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,061 filed January 31, 2000 which claimed the benefit of priority of provisional Patent Application

Serial No. 60/119,406 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the drug mixture of the invention may be delivered via a coated hydrogel controlled release dosage form having at least three components: (a) a composition containing the GPI, (b) a water-swellaable composition wherein the water-swellaable composition is in a separate region within a core formed by the drug-containing composition and the water-swellaable composition, and (c) a coating around the core that is water-permeable, water-insoluble, and has a least one delivery port therethrough. In use, the core imbibes water through the coating, swelling the water-swellaable composition and increasing the pressure within the core, and fluidizing the GPI-containing composition. Because the coating remains intact, the GPI-containing composition is extruded out of the delivery port into an environment of use. The polymer may be delivered in a separate dosage form, may be included in the GPI-containing composition, may comprise a separate composition that occupies a separate region within the core, or may constitute all or part of a coating applied to the dosage form. Examples of such dosage forms are more fully disclosed in commonly assigned pending Provisional Application Serial No. 60/171,968 filed December 23, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions may be administered as multiparticulates. Multiparticulates generally refer to dosage forms that comprise a multiplicity of particles that may range in size from about 10  $\mu\text{m}$  to about 2 mm, more typically about 100  $\mu\text{m}$  to 1 mm in diameter. Such multiparticulates may be packaged, for example, in a capsule such as a gelatin capsule or a capsule formed from an aqueous-soluble polymer such as HPMCAS, HPMC or starch or they may be

dosed as a suspension or slurry in a liquid. Such particulates may be made by any known process such as wet and dry granulation processes or melt congeal processes such as those previously described for forming amorphous  
5 GPI. For example, the GPI and a glyceride such as hydrogenated vegetable oil, a vegetable or synthetic fat or a wax such as paraffin may be blended and fed to a melt congeal process as a solid or liquid, followed by cooling to form beads comprised of amorphous GPI and the  
10 excipient.

The so-formed beads may then be blended with one or more concentration-enhancing polymers with or without additional excipients to form a multiparticulate dosage form. Alternatively, a high melting point  
15 concentration-enhancing polymer such as HPMCAS may be blended with the GPI and the fat or wax fed as a solid blend to a melt congeal process or the blend may be heated such that the GPI and the fat or wax melt to form a slurry of concentration-enhancing polymer particles in  
20 molten GPI and fat or wax. The resulting material comprises beads or particles consisting of an amorphous dispersion of GPI in the fat or wax with concentration-enhancing polymer particles trapped therein. Alternatively, a dispersion of the GPI in a  
25 concentration-enhancing polymer may be blended with a fat or wax and then fed to a melt congeal process as a solid or a slurry of the dispersion in the molten fat or wax. Such processing yields particles or beads consisting of particles of dispersion trapped in the solidified fat or  
30 wax matrix.

Similar multiparticulate dosage forms may be made with the various compositions of this invention but using excipients suited to the bead-forming or granule-forming process chosen. For example, when granules are  
35 formed by extrusion/spheronization processes the dispersion or other composition may be blended with, for

example, microcrystalline cellulose or other cellulosic polymer to aid in processing.

In any case, the resulting particles may themselves constitute the multiparticulate dosage form or  
5 they may be coated by various film-forming materials such as enteric polymers or water-swellaable or water-soluble polymers, or they may be combined with other excipients or vehicles to aid in dosing to patients.

Alternatively, the compositions of the present  
10 invention may be co-administered, meaning that the GPI can be administered separately from, but within the same general time frame as, the polymer. Thus, amorphous GPI can, for example, be administered in its own dosage form which is taken at approximately the same time as the  
15 polymer which is in a separate dosage form. If administered separately, it is generally preferred to administer both the GPI and the polymer within 60 minutes, more preferably within 15 minutes, of each other, so that the two are present together in the  
20 environment of use. When not administered simultaneously, the polymer is preferably administered prior to the amorphous GPI.

In addition to the above additives or  
excipients, use of any conventional materials and  
25 procedures for preparation of suitable dosage forms using the compositions of this invention known by those skilled in the art are potentially useful.

In another aspect, the present invention  
concerns the treatment of diabetes, including impaired  
30 glucose tolerance, insulin resistance, insulin dependent diabetes mellitus (Type 1) and non-insulin dependent diabetes mellitus (NIDDM or Type 2). Also included in the treatment of diabetes are the treatment of the diabetic complications, such as neuropathy, nephropathy,  
35 retinopathy or cataracts. The compositions of the present invention can also be used for diabetes prevention.

Diabetes can be treated by administering to a patient having diabetes (Type 1 or Type 2), insulin resistance, impaired glucose tolerance, or any of the diabetic complications such as neuropathy, nephropathy, retinopathy or cataracts, a therapeutically effective amount of a composition of the present invention. It is also contemplated that diabetes be treated by administering a composition of the present invention in combination with other agents that can be used to treat diabetes.

Representative agents that can be used to treat diabetes include insulin and insulin analogs (e.g. LysPro insulin); GLP-1 (7-37) (insulinotropin) and GLP-1 (7-36)-NH<sub>2</sub>; sulfonylureas and analogs: chlorpropamide, glibenclamide, tolbutamide, tolazamide, acetohexamide, glypizide, glimepiride, repaglinide, meglitinide; biguanides: metformin, phenformin, buformin;  $\alpha$ -antagonists and imidazolines: midaglizole, isaglidole, deriglidole, idazoxan, efaroxxan, fluparoxan; Other insulin secretagogues: linoglitazone, A-4166; glitazones: ciglitazone, pioglitazone, englitazone, troglitazone, darglitazone, rosiglitazone; PPAR-gamma agonists; fatty acid oxidation inhibitors: clomoxir, etomoxir;  $\alpha$ -glucosidase inhibitors: acarbose, miglitol, emiglitate, voglibose, MDL-25,637, camiglibose, MDL-73,945;  $\beta$ -agonists: BRL 35135, BRL 37344, Ro 16-8714, ICI D7114, CL 316,243; phosphodiesterase inhibitors: L-386,398; lipid-lowering agents: benfluorex; antiobesity agents: fenfluramine; vanadate and vanadium complexes (e.g. Naglivan®) and peroxovanadium complexes; amylin antagonists; glucagon antagonists; gluconeogenesis inhibitors; somatostatin analogs and antagonists; antilipolytic agents: nicotinic acid, acipimox, WAG 994. Any combination of agents can be administered as described above.

In addition to the categories and compounds mentioned above, the compositions of the present invention can be administered in combination with thyromimetic compounds, aldose reductase inhibitors, glucocorticoid receptor antagonists, NHE-1 inhibitors, or sorbitol dehydrogenase inhibitors, or combinations thereof, to treat or prevent diabetes, insulin resistance, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia, particularly myocardial ischemia.

It is generally accepted that thyroid hormones, specifically, biologically active iodothyronines, are critical to normal development and to maintaining metabolic homeostasis. Thyroid hormones stimulate the metabolism of cholesterol to bile acids and enhance the lipolytic responses of fat cells to other hormones. U.S. Patent Numbers 4,766,121; 4,826,876; 4,910,305; and 5,061,798 disclose certain thyroid hormone mimetics (thyromimetics), namely, 3,5-dibromo-3'-[6-oxo-3(1H)-pyridazinylmethyl]-thyronines. U.S. Patent Number 5,284,971 discloses certain thyromimetic cholesterol lowering agents, namely, 4-(3-cyclohexyl-4-hydroxy or -methoxy phenylsulfonyl)-3,5 dibromo-phenylacetic compounds. U.S. Patent Numbers 5,401,772; 5,654,468; and 5,569,674 disclose certain thyromimetics that are lipid lowering agents, namely, heteroacetic acid derivatives. In addition, certain oxamic acid derivatives of thyroid hormones are known in the art. For example, N. Yokoyama, et al. in an article published in the Journal of Medicinal Chemistry, 38 (4): 695-707 (1995) describe replacing a -CH<sub>2</sub> group in a naturally occurring metabolite of T<sub>3</sub> with an -NH group resulting in -HNCOCO<sub>2</sub>H. Likewise, R.E. Steele et al. in an article published in International Congressional Service (Atherosclerosis X) 1066: 321-324 (1995) and Z.F. Stephan et al. in an

article published in Atherosclerosis, 126: 53-63 (1996), describe certain oxamic acid derivatives useful as lipid-lowering thyromimetic agents, yet devoid of undesirable cardiac activities.

5           Each of the thyromimetic compounds referenced above and other thyromimetic compounds can be used in combination with the compositions of the present invention to treat or prevent diabetes, insulin resistance, diabetic neuropathy, diabetic nephropathy,  
10   diabetic retinopathy, cataracts, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia.

          The compositions of the present invention can also be used in combination with aldose reductase  
15   inhibitors. Aldose reductase inhibitors constitute a class of compounds that have become widely known for their utility in preventing and treating conditions arising from complications of diabetes, such as diabetic neuropathy and nephropathy. Such compounds are well  
20   known to those skilled in the art and are readily identified by standard biological tests. For example, the aldose reductase inhibitors zopolrestat, 1-phthalazineacetic acid, 3,4-dihydro-4-oxo-3-[[5-(trifluoromethyl)-2-benzothiazolyl]methyl]-, and related  
25   compounds are described in U.S. patent 4,939,140 to Larson et al.

          Aldose reductase inhibitors have been taught for use in lowering lipid levels in mammals. See, for example, U. S. patent 4,492,706 to Kallai-sanfacon and EP  
30   0 310 931 A2 (Ethyl Corporation).

          U. S. patent 5,064,830 to Going discloses the use of certain oxophthalazinyll acetic acid aldose reductase inhibitors, including zopolrestat, for lowering of blood uric acid levels.

35           Commonly assigned U.S. patent 5,391,551 discloses the use of certain aldose reductase inhibitors, including zopolrestat, for lowering blood lipid levels in

humans. The disclosure teaches that therapeutic utilities derive from the treatment of diseases caused by an increased level of triglycerides in the blood, such diseases include cardiovascular disorders such as thrombosis, arteriosclerosis, myocardial infarction, and angina pectoris. A preferred aldose reductase inhibitor is 1-phthalazineacetic acid, 3,4-dihydro-4-oxo-3-[[5-trifluoromethyl)-2-benzothiazolyl]methyl]-, also known as zopolrestat.

The term aldose reductase inhibitor refers to compounds that inhibit the bioconversion of glucose to sorbitol, which is catalyzed by the enzyme aldose reductase. Any aldose reductase inhibitor may be used in a combination with a composition of the present invention. Aldose reductase inhibition is readily determined by those skilled in the art according to standard assays (J. Malone, Diabetes, 29:861-864 (1980); "Red Cell Sorbitol, an Indicator of Diabetic Control"). A variety of aldose reductase inhibitors are described herein; however, other aldose reductase inhibitors useful in the compositions and methods of this invention will be known to those skilled in the art. The bioconversion of glucose to sorbitol. The activity of an aldose reductase inhibitor in a tissue can be determined by testing the amount of aldose reductase inhibitor that is required to lower tissue sorbitol (i.e., by inhibiting the further production of sorbitol consequent to blocking aldose reductase) or lower tissue fructose (by inhibiting the production of sorbitol consequent to blocking aldose reductase and consequently the production of fructose). Accordingly, examples of aldose reductase inhibitors useful in the compositions, combinations and methods of the present invention include:

1. 3-(4-bromo-2-fluorobenzyl)-3,4-dihydro-4-oxo-1-phthalazineacetic acid (ponalrestat, US 4,251,528);




2. N[[ (5-trifluoromethyl)-6-methoxy-1-naphthalenyl]thioxomethyl]-N-methylglycine (tolrestat, US 4,600,724);
3. 5-[(Z,E)- $\beta$ -methylcinnamylidene]-4-oxo-2-thioxo-3-thiazolideneacetic acid (epalrestat, US 4,464,382, US 4,791,126, US 4,831,045);
4. 3-(4-bromo-2-fluorobenzyl)-7-chloro-3,4-dihydro-2,4-dioxo-1(2H)-quinazolineacetic acid (zenarestat, US 4,734,419, and 4,883,800);
5. 2R,4R-6,7-dichloro-4-hydroxy-2-methylchroman-4-acetic acid (US 4,883,410);
6. 2R,4R-6,7-dichloro-6-fluoro-4-hydroxy-2-methylchroman-4-acetic acid (US 4,883,410);
7. 3,4-dihydro-2,8-diisopropyl-3-oxo-2H-1,4-benzoxazine-4-acetic acid (US 4,771,050);
8. 3,4-dihydro-3-oxo-4-[(4,5,7-trifluoro-2-benzothiazolyl)methyl]-2H-1,4-benzothiazine-2-acetic acid (SPR-210, U.S. 5,252,572);
9. N-[3,5-dimethyl-4-[(nitromethyl)sulfonyl]phenyl]-2-methyl-benzeneacetamide (ZD5522, U.S. 5,270,342 and U.S. 5,430,060);
10. (S)-6-fluorospiro[chroman-4,4'-imidazolidine]-2,5'-dione (sorbiniol, US 4,130,714);
11. d-2-methyl-6-fluoro-spiro(chroman-4',4'-imidazolidine)-2',5'-dione (US 4,540,704);
12. 2-fluoro-spiro(9H-fluorene-9,4'-imidazolidine)2',5'-dione (US 4,438,272);
13. 2,7-di-fluoro-spiro(9H-fluorene-9,4'-imidazolidine)2',5'-dione (US 4,436,745, US 4,438,272);
14. 2,7-di-fluoro-5-methoxy-spiro(9H-fluorene-9,4'-imidazolidine)2',5'-dione (US 4,436,745, US 4,438,272);
15. 7-fluoro-spiro(5H-indenol[1,2-b]pyridine-5,3'-pyrrolidine)2,5'-dione (US 4,436,745, US 4,438,272);
16. d-cis-6'-chloro-2',3'-dihydro-2'-methyl-spiro-(imidazolidine-4,4'-4'-H-pyrano(2,3-b)pyridine)-2,5-dione (US 4,980,357);

17. spiro[imidazolidine-4,5' (6H) -  
quinoline]2,5-dione-3'-chloro-7,'8'-dihydro-7'-methyl-  
(5'-cis) (US 5,066,659);

18. (2S,4S)-6-fluoro-2',5'-dioxospiro(chroman-  
5 4,4'-imidazolidine)-2-carboxamide (US 5,447,946); and

19. 2-[(4-bromo-2-fluorophenyl)methyl]-6-  
fluorospiro[isoquinoline-4(1H),3'-pyrrolidine]-  
1,2',3,5'(2H)-tetrone (ARI-509, US 5,037,831).

Other aldose reductase inhibitors include  
10 compounds having Formula Ia below



20 or a pharmaceutically acceptable salt or prodrug thereof,  
wherein the substituents of Formula Ia are as follows:

Z is O or S;

R<sup>1</sup> is hydroxy or a group capable of being  
removed in vivo to produce a compound of Formula I  
25 wherein R<sup>1</sup> is OH; and

X and Y are the same or different and are  
selected from hydrogen, trifluoromethyl, fluoro, and  
chloro.

A preferred subgroup within the above group of  
30 aldose reductase inhibitors includes numbered compounds  
1, 2, 3, 4, 5, 6, 9, 10, and 17, and the following  
compounds of Formula Ia:

20. 3,4-dihydro-3-(5-fluorobenzothiazol-2-  
ylmethyl)-4-oxophthalazin-1-yl-acetic acid [R<sup>1</sup>=hydroxy;  
35 X=F; Y=H];

21. 3-(5,7-difluorobenzothiazol-2-ylmethyl)-  
3,4-dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy;  
X=Y=F];
22. 3-(5-chlorobenzothiazol-2-ylmethyl)-3,4-  
5 dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy;  
X=Cl; Y=H];
23. 3-(5,7-dichlorobenzothiazol-2-ylmethyl)-  
3,4-dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy;  
X=Y=Cl];
- 10 24. 3,4-dihydro-4-oxo-3-(5-  
trifluoromethylbenzoxazol-2-ylmethyl)phthalazin-1-  
ylacetic acid [ $R^1$ =hydroxy; X=CF<sub>3</sub>; Y=H];
25. 3,4-dihydro-3-(5-fluorobenzoxazol-2-  
ylmethyl)-4-oxophthalazin-1-yl-acetic acid [ $R^1$ =hydroxy;  
15 X=F; Y=H];
26. 3-(5,7-difluorobenzoxazol-2-ylmethyl)-3,4-  
dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy;  
X=Y=F];
27. 3-(5-chlorobenzoxazol-2-ylmethyl)-3,4-  
20 dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy; X=Cl;  
Y=H];
28. 3-(5,7-dichlorobenzoxazol-2-ylmethyl)-3,4-  
dihydro-4-oxophthalazin-1-ylacetic acid [ $R^1$ =hydroxy;  
X=Y=Cl]; and
- 25 29. zopolrestat; 1-phthalazineacetic acid,  
3,4-dihydro-4-oxo-3-[[5-(trifluoromethyl)-2-  
benzothiazolyl]methyl]- [ $R^1$ =hydroxy; X=trifluoromethyl;  
Y=H].

30 In compounds 20-23, and 29 Z is S. In compounds  
24-28, Z is O.


Of the above subgroup, compounds 20-29 are more  
preferred with 29 especially preferred. Procedures for  
making the aldose reductase inhibitors of formula Ia can  
be found in PCT publication number WO 99/26659.

35 Each of the aldose reductase inhibitors  
referenced above and other aldose reductase inhibitors  
can be used in combination with the compounds of the

present invention to treat diabetes, insulin resistance, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia.

The compositions of the present invention can also be used in combination with glucocorticoid receptor antagonists. The glucocorticoid receptor (GR) is present in glucocorticoid responsive cells where it resides in the cytosol in an inactive state until it is stimulated by an agonist. Upon stimulation the glucocorticoid receptor translocates to the cell nucleus where it specifically interacts with DNA and/or protein(s) and regulates transcription in a glucocorticoid responsive manner. Two examples of proteins that interact with the glucocorticoid receptor are the transcription factors, API and NFκ-B. Such interactions result in inhibition of API- and NFκ-B- mediated transcription and are believed to be responsible for the anti-inflammatory activity of endogenously administered glucocorticoids. In addition, glucocorticoids may also exert physiologic effects independent of nuclear transcription. Biologically relevant glucocorticoid receptor agonists include cortisol and corticosterone. Many synthetic glucocorticoid receptor agonists exist including dexamethasone, prednisone and prednisilone. By definition, glucocorticoid receptor antagonists bind to the receptor and prevent glucocorticoid receptor agonists from binding and eliciting GR mediated events, including transcription. RU486 is an example of a non-selective glucocorticoid receptor antagonist. GR antagonists can be used in the treatment of diseases associated with an excess or a deficiency of glucocorticoids in the body. As such, they may be used to treat the following: obesity, diabetes, cardiovascular disease, hypertension, Syndrome X, depression, anxiety, glaucoma, human immunodeficiency virus (HIV) or acquired immunodeficiency

syndrome (AIDS), neurodegeneration (for example, Alzheimer's and Parkinson's), cognition enhancement, Cushing's Syndrome, Addison's Disease, osteoporosis, frailty, inflammatory diseases (such as osteoarthritis, rheumatoid arthritis, asthma and rhinitis), tests of adrenal function, viral infection, immunodeficiency, immunomodulation, autoimmune diseases, allergies, wound healing, compulsive behavior, multi-drug resistance, addiction, psychosis, anorexia, cachexia, post-traumatic stress syndrome, post-surgical bone fracture, medical catabolism and prevention of muscle frailty. Examples of GR antagonists that can be used in combination with a compound of the present invention include compounds of Formula Ib below:




an isomer thereof, a prodrug of said compound or isomer, or a pharmaceutically acceptable salt of said compound, isomer or prodrug wherein the substituents of Formula Ib are as follows:


$m$  is 1 or 2;

- - - represents an optional bond;

A is selected from the group consisting of



and



A-5

D is CR<sub>7</sub>, CR<sub>7</sub>R<sub>16</sub>, N, NR<sub>7</sub> or O;

E is C, CR<sub>6</sub> or N;

F is CR<sub>4</sub>, CR<sub>4</sub>R<sub>5</sub> or O;

G, H and I together with 2 carbon atoms from the A-ring or 2 carbon atoms from the B-ring form a 5-membered heterocyclic ring comprising one or more N, O or S atoms; provided that there is at most one of O and S per ring;

J, K, L and M together with 2 carbon atoms from the B-ring forms a 6-membered heterocyclic ring comprising 1 or more N atoms;

X is a) absent, b) -CH<sub>2</sub>-, c) -CH(OH)- or d) -C(O)-;

R<sub>1</sub> is a) -H, b) -Z-CF<sub>3</sub>, c) -(C<sub>1</sub>-C<sub>6</sub>)alkyl, d) -(C<sub>2</sub>-C<sub>6</sub>)alkenyl, e) -(C<sub>2</sub>-C<sub>6</sub>)alkynyl, f) -CHO, g) -CH=N-OR<sub>12</sub>, h) -Z-C(O)OR<sub>12</sub>, i) -Z-C(O)-NR<sub>12</sub>R<sub>13</sub>, j) -Z-C(O)-NR<sub>12</sub>-Z-het, k) -Z-NR<sub>12</sub>R<sub>13</sub>, l) -Z-NR<sub>12</sub>het, m) -Z-het, n) -Z-O-het, o) -Z-aryl', p) -Z-O-aryl', q) -CHOH-aryl' or r) -C(O)-aryl' wherein aryl' in substituents o) to r) is substituted independently with 0, 1 or 2 of the following: -Z-OH, -Z-NR<sub>12</sub>R<sub>13</sub>, -Z-NR<sub>12</sub>-het, -C(O)NR<sub>12</sub>R<sub>13</sub>, -C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -C(O)OH, -C(O)-het, -NR<sub>12</sub>-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, -NR<sub>12</sub>-C(O)-(C<sub>2</sub>-C<sub>6</sub>)alkenyl, -NR<sub>12</sub>-C(O)-(C<sub>2</sub>-C<sub>6</sub>)alkynyl, -NR<sub>12</sub>-C(O)-Z-het, -CN, -Z-het, -O-(C<sub>1</sub>-C<sub>3</sub>)alkyl-C(O)-NR<sub>12</sub>R<sub>13</sub>, -O-(C<sub>1</sub>-C<sub>3</sub>)alkyl-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -NR<sub>12</sub>-Z-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -N(Z-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>, -NR<sub>12</sub>-Z-C(O)-NR<sub>12</sub>R<sub>13</sub>, -Z-NR<sub>12</sub>-SO<sub>2</sub>-R<sub>13</sub>, -NR<sub>12</sub>-SO<sub>2</sub>-het, -C(O)H, -Z-NR<sub>12</sub>-Z-O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -Z-NR<sub>12</sub>-Z-NR<sub>12</sub>R<sub>13</sub>, -Z-NR<sub>12</sub>-(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, -Z-N(Z-O(C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>, -SO<sub>2</sub>R<sub>12</sub>,

-SOR<sub>12</sub>, -SR<sub>12</sub>, -SO<sub>2</sub>NR<sub>12</sub>R<sub>13</sub>, -O-C(O)-(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
-O-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>4</sub>)alkyl, -halo or -CF<sub>3</sub>;

Z for each occurrence is independently a)  
-(C<sub>0</sub>-C<sub>6</sub>)alkyl, b) -(C<sub>2</sub>-C<sub>6</sub>)alkenyl or c) -(C<sub>2</sub>-C<sub>6</sub>)alkynyl;

5 R<sub>2</sub> is a) -H, b) -halo, c) -OH, d) -(C<sub>1</sub>-C<sub>6</sub>)alkyl  
substituted with 0 or 1 -OH, e) -NR<sub>12</sub>R<sub>13</sub>, f)  
-Z-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, g) -Z-C(O)NR<sub>12</sub>R<sub>13</sub>, h) -O-(C<sub>1</sub>-C<sub>6</sub>)alkyl,  
i) -Z-O-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, j)  
-Z-O-(C<sub>1</sub>-C<sub>3</sub>)alkyl-C(O)-NR<sub>12</sub>R<sub>13</sub>, k)  
10 -Z-O-(C<sub>1</sub>-C<sub>3</sub>)alkyl-C(O)-O(C<sub>1</sub>-C<sub>6</sub>)alkyl, l) -O-(C<sub>2</sub>-C<sub>6</sub>)alkenyl,  
m) -O-(C<sub>2</sub>-C<sub>6</sub>)alkynyl, n) -O-Z-het, o) -COOH, p) -C(OH)R<sub>12</sub>R<sub>13</sub>  
or q) -Z-CN;

R<sub>3</sub> is a) -H, b) -(C<sub>1</sub>-C<sub>10</sub>)alkyl wherein 1 or 2  
carbon atoms, other than the connecting carbon atom, may  
15 optionally be replaced with 1 or 2 heteroatoms  
independently selected from S, O and N and wherein each  
carbon atom is substituted with 0, 1 or 2 R<sub>y</sub>, c)  
-(C<sub>2</sub>-C<sub>10</sub>)alkenyl substituted with 0, 1 or 2 R<sub>y</sub>, d)  
-(C<sub>2</sub>-C<sub>10</sub>)alkynyl wherein 1 carbon atom, other than the  
20 connecting carbon atom, may optionally be replaced with 1  
oxygen atom and wherein each carbon atom is substituted  
with 0, 1 or 2 R<sub>y</sub>, e) -CH=C=CH<sub>2</sub>, f) -CN, g)  
-(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, h) -Z-aryl, i) -Z-het, j)  
-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, k) -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, l) -Z-S-R<sub>12</sub>, m)  
25 -Z-S(O)-R<sub>12</sub>, n) -Z-S(O)<sub>2</sub>-R<sub>12</sub>, o) -CF<sub>3</sub>, p) -NR<sub>12</sub>O-(C<sub>1</sub>-C<sub>6</sub>)alkyl  
or q) -CH<sub>2</sub>OR<sub>y</sub>;

provided that one of R<sub>2</sub> and R<sub>3</sub> is absent when  
there is a double bond between CR<sub>2</sub>R<sub>3</sub> (the 7 position) and  
the F moiety (the 8 position) of the C-ring;

30 R<sub>y</sub> for each occurrence is independently a) -OH,  
b) -halo, c) -Z-CF<sub>3</sub>, d) -Z-CF(C<sub>1</sub>-C<sub>3</sub>alkyl)<sub>2</sub>, e) -CN, f)  
-NR<sub>12</sub>R<sub>13</sub>, g) -(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, h) -(C<sub>3</sub>-C<sub>6</sub>)cycloalkenyl, i)  
-(C<sub>0</sub>-C<sub>3</sub>)alkyl-aryl, j) -het or k) -N<sub>3</sub>;

or R<sub>2</sub> and R<sub>3</sub> are taken together to form a)  
35 =CHR<sub>11</sub>, b) =NOR<sub>11</sub>, c) =O, d) =N-NR<sub>12</sub>, e) =N-NR<sub>12</sub>-C(O)-R<sub>12</sub>, f)  
oxiranyl or g) 1,3-dioxolan-4-yl;

$R_4$  and  $R_5$  for each occurrence are independently  
 a) -H, b) -CN, c)  $-(C_1-C_6)$ alkyl substituted with 0 to 3  
 halo, d)  $-(C_2-C_6)$ alkenyl substituted with 0 to 3 halo, e)  
 $-(C_2-C_6)$ alkynyl substituted with 0 to 3 halo, f)  
 5 -O- $(C_1-C_6)$ alkyl substituted with 0 to 3 halo, g)  
 -O- $(C_2-C_6)$ alkenyl substituted with 0 to 3 halo, h)  
 -O- $(C_2-C_6)$ alkynyl substituted with 0 to 3 halo, i) halo,  
 j) -OH, k)  $(C_3-C_6)$ cycloalkyl or l)  $(C_3-C_6)$ cycloalkenyl;

or  $R_4$  and  $R_5$  are taken together to form =O;

10  $R_6$  is a) -H, b) -CN, c)  $-(C_1-C_6)$ alkyl  
 substituted with 0 to 3 halo, d)  $-(C_2-C_6)$ alkenyl  
 substituted with 0 to 3 halo, e)  $-(C_2-C_6)$ alkynyl  
 substituted with 0 to 3 halo or f) -OH;

$R_7$  and  $R_{16}$  for each occurrence are independently  
 15 a) -H, b) -halo, c) -CN, d)  $-(C_1-C_6)$ alkyl substituted with  
 0 to 3 halo, e)  $-(C_2-C_6)$ alkenyl substituted with 0 to 3  
 halo or f)  $-(C_2-C_6)$ alkynyl substituted with 0 to 3 halo;  
 provided that  $R_7$  is other than -CN or -halo when D is  $NR_7$ ;

or  $R_7$  and  $R_{16}$  are taken together to form =O;

20  $R_8$ ,  $R_9$ ,  $R_{14}$  and  $R_{15}$  for each occurrence are  
 independently a) -H, b) -halo, c)  $(C_1-C_6)$ alkyl substituted  
 with 0 to 3 halo, d)  $-(C_2-C_6)$ alkenyl substituted with 0 to  
 3 halo, e)  $-(C_2-C_6)$ alkynyl substituted with 0 to 3 halo,  
 f) -CN, g)  $-(C_3-C_6)$ cycloalkyl, h)  $-(C_3-C_6)$ cycloalkenyl, i)  
 25 -OH, j) -O- $(C_1-C_6)$ alkyl, k) -O- $(C_1-C_6)$ alkenyl, l)  
 -O- $(C_1-C_6)$ alkynyl, m)  $-NR_{12}R_{13}$ , n)  $-C(O)OR_{12}$  or o)  
 $-C(O)NR_{12}R_{13}$ ;

or  $R_8$  and  $R_9$  are taken together on the C-ring to  
 form =O; provided that when m is 2, only one set of  $R_8$  and  
 30  $R_9$  are taken together to form =O;

or  $R_{14}$  and  $R_{15}$  are taken together to form =O;  
 provided that when  $R_{14}$  and  $R_{15}$  are taken together to form  
 =O, D is other than  $CR_7$  and E is other than C;

$R_{10}$  is a)  $-(C_1-C_{10})$ alkyl substituted with 0 to 3  
 35 substituents independently selected from -halo, -OH and  
 $-N_3$ , b)  $-(C_2-C_{10})$ alkenyl substituted with 0 to 3  
 substituents independently selected from -halo, -OH and



- N<sub>3</sub>, c) -(C<sub>2</sub>-C<sub>10</sub>)alkynyl substituted with 0 to 3 substituents independently selected from -halo, -OH and -N<sub>3</sub>, d) -halo, e) -Z-CN, f) -OH, g) -Z-het, h) -Z-NR<sub>12</sub>R<sub>13</sub>, i) -Z-C(O)-het, j) -Z-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, k) -Z-C(O)-NR<sub>12</sub>R<sub>13</sub>,  
 5 l) -Z-C(O)-NR<sub>12</sub>-Z-CN, m) -Z-C(O)-NR<sub>12</sub>-Z-het, n) -Z-C(O)-NR<sub>12</sub>-Z-aryl, o) -Z-C(O)-NR<sub>12</sub>-Z-NR<sub>12</sub>R<sub>13</sub>, p) -Z-C(O)-NR<sub>12</sub>-Z-O(C<sub>1</sub>-C<sub>6</sub>)alkyl, q) -(C<sub>1</sub>-C<sub>6</sub>)alkyl-C(O)OH, r) -Z-C(O)O(C<sub>1</sub>-C<sub>6</sub>)alkyl, s) -Z-O-(C<sub>0</sub>-C<sub>6</sub>)alkyl-het, t) -Z-O-(C<sub>0</sub>-C<sub>6</sub>)alkyl-aryl, u) -Z-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl substituted  
 10 with 0 to 2 R<sub>x</sub>, v) -Z-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl-CH(O), w) -Z-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl-NR<sub>12</sub>-het, x) -Z-O-Z-het-Z-het, y) -Z-O-Z-het-Z-NR<sub>12</sub>R<sub>13</sub>, z) -Z-O-Z-het-C(O)-het, a1) -Z-O-Z-C(O)-het, b1) -Z-O-Z-C(O)-het-het, c1) -Z-O-Z-C(O)-(C<sub>1</sub>-C<sub>6</sub>)alkyl, d1) -Z-O-Z-C(S)-NR<sub>12</sub>R<sub>13</sub>, e1)  
 15 -Z-O-Z-C(O)-NR<sub>12</sub>R<sub>13</sub>, f1) -Z-O-Z-(C<sub>1</sub>-C<sub>3</sub>)alkyl-C(O)-NR<sub>12</sub>R<sub>13</sub>, g1) -Z-O-Z-C(O)-O(C<sub>1</sub>-C<sub>6</sub>)alkyl, h1) -Z-O-Z-C(O)-OH, i1) -Z-O-Z-C(O)-NR<sub>12</sub>-O(C<sub>1</sub>-C<sub>6</sub>)alkyl, j1) -Z-O-Z-C(O)-NR<sub>12</sub>-OH, k1) -Z-O-Z-C(O)-NR<sub>12</sub>-Z-NR<sub>12</sub>R<sub>13</sub>, l1) -Z-O-Z-C(O)-NR<sub>12</sub>-Z-het, m1) -Z-O-Z-C(O)-NR<sub>12</sub>-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub>)alkyl, n1)  
 20 -Z-O-Z-C(=NR<sub>12</sub>)(NR<sub>12</sub>R<sub>13</sub>), o1) -Z-O-Z-C(=NOR<sub>12</sub>)(NR<sub>12</sub>R<sub>13</sub>), p1) -Z-NR<sub>12</sub>-C(O)-O-Z-NR<sub>12</sub>R<sub>13</sub>, q1) -Z-S-C(O)-NR<sub>12</sub>R<sub>13</sub>, r1) -Z-O-SO<sub>2</sub>-(C<sub>1</sub>-C<sub>6</sub>)alkyl, s1) -Z-O-SO<sub>2</sub>-aryl, t1) -Z-O-SO<sub>2</sub>-NR<sub>12</sub>R<sub>13</sub>, u1) -Z-O-SO<sub>2</sub>-CF<sub>3</sub>, v1) -Z-NR<sub>12</sub>C(O)OR<sub>13</sub> or w1) -Z-NR<sub>12</sub>C(O)R<sub>13</sub>;  
 25 or R<sub>9</sub> and R<sub>10</sub> are taken together on the moiety of formula A-5 to form a) = O or b) = NOR<sub>12</sub>;  
 R<sub>11</sub> is a) -H, b) -(C<sub>1</sub>-C<sub>5</sub>)alkyl, c) -(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl or d) -(C<sub>0</sub>-C<sub>3</sub>)alkyl-aryl;  
 R<sub>12</sub> and R<sub>13</sub> for each occurrence are each  
 30 independently a) -H, b) -(C<sub>1</sub>-C<sub>6</sub>)alkyl wherein 1 or 2 carbon atoms, other than the connecting carbon atom, may optionally be replaced with 1 or 2 heteroatoms independently selected from S, O and N and wherein each carbon atom is substituted with 0 to 6 halo, c)  
 35 -(C<sub>2</sub>-C<sub>6</sub>)alkenyl substituted with 0 to 6 halo or d) -(C<sub>1</sub>-C<sub>6</sub>)alkynyl wherein 1 carbon atom, other than the connecting carbon atom, may optionally be replaced with 1

oxygen atom and wherein each carbon atom is substituted with 0 to 6 halo;

or  $R_{12}$  and  $R_{13}$  are taken together with N to form het;

5 or  $R_6$  and  $R_{14}$  or  $R_{15}$  are taken together to form 1,3-dioxolanyl;

aryl is a) phenyl substituted with 0 to 3  $R_x$ , b) naphthyl substituted with 0 to 3  $R_x$  or c) biphenyl substituted with 0 to 3  $R_x$ ;

10 het is a 5-, 6- or 7-membered saturated, partially saturated or unsaturated ring containing from one (1) to three (3) heteroatoms independently selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the  
15 above heterocyclic rings is fused to a benzene ring or another heterocycle; and the nitrogen may be in the oxidized state giving the N-oxide form; and substituted with 0 to 3  $R_x$ ;

$R_x$  for each occurrence is independently a)  
20 -halo, b) -OH, c)  $-(C_1-C_6)$ alkyl, d)  $-(C_2-C_6)$ alkenyl, e)  $-(C_2-C_6)$ alkynyl, f)  $-O(C_1-C_6)$ alkyl, g)  $-O(C_2-C_6)$ alkenyl, h)  $-O(C_2-C_6)$ alkynyl, i)  $-(C_0-C_6)$ alkyl- $NR_{12}R_{13}$ , j)  $-C(O)-NR_{12}R_{13}$ , k)  $-Z-SO_2R_{12}$ , l)  $-Z-SOR_{12}$ , m)  $-Z-SR_{12}$ , n)  $-NR_{12}-SO_2R_{13}$ , o)  $-NR_{12}-C(O)-R_{13}$ , p)  $-NR_{12}-OR_{13}$ , q)  $-SO_2-NR_{12}R_{13}$ , r) -CN, s)  
25  $-CF_3$ , t)  $-C(O)(C_1-C_6)$ alkyl, u) =O, v)  $-Z-SO_2$ -phenyl or w)  $-Z-SO_2$ -het';

aryl' is phenyl, naphthyl or biphenyl;

het' is a 5-, 6- or 7-membered saturated, partially saturated or unsaturated ring containing from  
30 one (1) to three (3) heteroatoms independently selected from the group consisting of nitrogen, oxygen and sulfur; and including any bicyclic group in which any of the above heterocyclic rings is fused to a benzene ring or another heterocycle;

35 provided that:

(1)  $X-R_1$  is other than hydrogen or methyl;

(2) when  $R_9$  and  $R_{10}$  are substituents on the A-ring, they are other than mono- or di-methoxy;

(3) when  $R_2$  and  $R_3$  are taken together to form  $=CHR_{11}$  or  $=O$  wherein  $R_{11}$  is  $-O(C_1-C_6)\text{alkyl}$ , then  $-X-R_1$  is  
5 other than  $(C_1-C_4)\text{alkyl}$ ;

(4) when  $R_2$  and  $R_3$  taken together are  $C=O$  and  $R_9$  is hydrogen on the A-ring; or when  $R_2$  is hydroxy,  $R_3$  is hydrogen and  $R_9$  is hydrogen on the A-ring, then  $R_{10}$  is other than  $-O-(C_1-C_6)\text{alkyl}$  or  $-O-CH_2\text{-phenyl}$  at the  
10 2-position of the A-ring;

(5) when  $X-R_1$  is  $(C_1-C_4)\text{alkyl}$ ,  $(C_2-C_4)\text{alkenyl}$  or  $(C_2-C_4)\text{alkynyl}$ ,  $R_9$  and  $R_{10}$  are other than mono-hydroxy or  $=O$ , including the diol form thereof, when taken together; and

(6) when  $X$  is absent,  $R_1$  is other than a moiety  
15 containing a heteroatom independently selected from N, O or S directly attached to the juncture of the B-ring and the C-ring. (See U.S. Provisional Patent Application number 60/132,130.)


Each of the glucocorticoid receptor antagonists referenced above and other glucocorticoid receptor antagonists can be used in combination with the compounds of the present invention to treat or prevent diabetes, hyperglycemia, hypercholesterolemia, hypertension,  
20 hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia.

The compositions of the present invention can also be used in combination with sorbitol dehydrogenase inhibitors. Sorbitol dehydrogenase inhibitors lower  
30 fructose levels and have been used to treat or prevent diabetic complications such as neuropathy, retinopathy, nephropathy, cardiomyopathy, microangiopathy, and macroangiopathy. U.S. patent numbers 5,728,704 and 5,866,578 disclose compounds and a method for treating or  
35 preventing diabetic complications by inhibiting the enzyme sorbitol dehydrogenase.

Each of the sorbitol dehydrogenase inhibitors referenced above and other sorbitol dehydrogenase inhibitors can be used in combination with the compounds of the present invention to treat diabetes, insulin resistance, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia.

The compositions of the present invention can also be used in combination with sodium-hydrogen exchanger Type 1 (NHE-1) inhibitors. NHE-1 inhibitors can be used to reduce tissue damage resulting from ischemia. Of great concern is tissue damage that occurs as a result of ischemia in cardiac, brain, liver, kidney, lung, gut, skeletal muscle, spleen, pancreas, nerve, spinal cord, retina tissue, the vasculature, or intestinal tissue. NHE-1 inhibitors can also be administered to prevent perioperative myocardial ischemic injury.

Examples of NHE-1 inhibitors include a compound having the Formula Ic



Formula Ic

a prodrug thereof or a pharmaceutically acceptable salt of said compound or of said prodrug, wherein the substituents of Formula Ic are as follows:

Z is carbon connected and is a five-membered, diaza, diunsaturated ring having two contiguous nitrogens, said ring optionally mono-, di-, or tri-substituted with up to three substituents independently selected from R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup>;  
or

Z is carbon connected and is a five-membered, triaza, diunsaturated ring, said ring optionally mono- or di-substituted with up to two substituents independently selected from R<sup>4</sup> and R<sup>5</sup>;

5                wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> are each independently hydrogen, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, M or M(C<sub>1</sub>-C<sub>4</sub>)alkyl, any of  
10   said previous (C<sub>1</sub>-C<sub>4</sub>)alkyl moieties optionally having from one to nine fluorines; said (C<sub>1</sub>-C<sub>4</sub>)alkyl or (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl optionally mono-or di-substituted independently with hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl,  
15   (C<sub>1</sub>-C<sub>4</sub>)alkyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl; and said (C<sub>3</sub>-C<sub>4</sub>)cycloalkyl optionally having from one to seven fluorines;

              wherein M is a partially saturated, fully  
20   saturated or fully unsaturated five to eight membered ring optionally having one to three heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to  
25   six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

              said M is optionally substituted, on one ring if the moiety is monocyclic, or one or both rings if the  
30   moiety is bicyclic, on carbon or nitrogen with up to three substituents independently selected from R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup>, wherein one of R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> is optionally a partially saturated, fully saturated, or fully unsaturated three to seven membered ring optionally having one to three  
35   heteroatoms selected independently from oxygen, sulfur and nitrogen optionally substituted with (C<sub>1</sub>-C<sub>4</sub>)alkyl and additionally R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are optionally hydroxy, nitro,

halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, formyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkynyl or (C<sub>5</sub>-C<sub>7</sub>)cycloalkenyl,

wherein said (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>7</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino or (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> substituents are optionally mono- substituted independently with hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoyl, (C<sub>1</sub>-C<sub>4</sub>)alkanoylamino, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxycarbonylamino, sulfonamido, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonamido, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylcarbamoyl, cyano, thiol, nitro, (C<sub>1</sub>-C<sub>4</sub>)alkylthio, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfinyl, (C<sub>1</sub>-C<sub>4</sub>)alkylsulfonyl or mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylaminosulfonyl or optionally substituted with one to nine fluorines. (See PCT patent application number PCT/IB99/00206)

Each of the NHE-1 inhibitors referenced above and other NHE-1 inhibitors can be used in combination with the compositions of the present invention to treat or prevent diabetes, insulin resistance, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hyperglycemia, hypercholesterolemia, hypertension, hyperinsulinemia, hyperlipidemia, atherosclerosis, or tissue ischemia.

Other features and embodiments of the invention will become apparent from the following examples which are given for illustration of the invention rather than for limiting its intended scope.

## EXAMPLES

## Example 1

This example discloses preparation of an amorphous solid dispersion of the GPI 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R, 4S)-dihydroxy-pyrrolidin-1-yl)-3-oxy-propyl]-amide ("Drug 1"), which has a solubility in water of 60 to 80  $\mu\text{g/mL}$  and a solubility in MFD solution of 183  $\mu\text{g/mL}$ . A dispersion of 25 wt% Drug 1 and 75 wt% polymer was made by first mixing Drug 1 in the solvent acetone together with a "medium fine" (AQUOT-MF) grade of the cellulosic enteric polymer HPMCAS (manufactured by Shin Etsu) to form a solution. The solution comprised 1.25 wt% Drug 1, 3.75 wt% HPMCAS, and 95 wt% acetone. This solution was then spray-dried by directing an atomizing spray using a two-fluid external-mix spray nozzle at 2.6 bar (37 psig) at a feed rate of 175 to 180 g/min into the stainless-steel chamber of a Niro XP spray-dryer, maintained at a temperature of 180°C at the inlet and 69°C at the outlet.

The resulting amorphous solid spray-dried dispersion (SDD) was collected via a cyclone and then dried in a Gruenberg solvent tray-dryer by spreading the spray-dried particles onto polyethylene-lined trays to a depth of not more than 1 cm and then drying them at 40°C for at least 8 hours.

## Examples 2-7

Examples 2 through 7 were prepared using the same process as in Example 1, with the exception that different dispersion polymers and different amounts of drug and polymer were used. The variables are noted in Table 1. The SDD of Example 2 was prepared using the Niro PSD-1 spray-dryer. The SDDs of Examples 3-7 were prepared using a "mini" spray dryer, which consisted of an atomizer in the top cap of a vertically oriented stainless steel pipe. The atomizer was a two-fluid nozzle (Spraying Systems Co. 1650 fluid cap and 64 air

cap) where the atomizing gas was nitrogen delivered to the nozzle at 100°C and a flow of 15 gm/min, and the spray-dried solution was delivered to the nozzle at room temperature and a flow rate of 1 gm/min using a syringe pump. Filter paper with a supporting screen was clamped to the bottom end of the pipe to collect the solid spray-dried material and to allow the nitrogen and evaporated solvent to escape.



Table 1

Ex. No.	Drug	Drug Mass	Polymer*	Polymer Mass	Drug Conc. in the Dispersion (wt%)	Solv	Solv Mass	Spray Dryer
1	1	991g	HPMCAS-MF	3009g	25	acetone	101,670g	Niro XP
2	1	30g	HPMCAS-MF	30g	50	acetone	2,340g	Niro PSD-1
3	1	25mg	HPMC	75mg	25	acetone	10g	Mini
4	1	25mg	PVP	75mg	25	acetone	10g	Mini
5	1	25mg	CAP	75mg	25	acetone	10g	Mini
6	1	25mg	CAT	75mg	25	acetone	10g	Mini
7	1	25mg	HPMCP	75mg	25	acetone	10g	Mini

\* Polymer designations: HPMCAS = hydroxypropyl methyl cellulose acetate succinate, HPMC = hydroxypropyl methyl cellulose, PVP = polyvinylpyrrolidone, CAP = cellulose acetate phthalate, CAT = cellulose acetate trimellitate, HPMCP = hydroxypropyl methyl cellulose phthalate.

## Examples 8-9

Example 8 was prepared by rotoevaporating a polymer:drug solution to dryness. The solution consisted of 7.5 wt% Drug 1, 7.5 wt% HPMCAS-MF, 80.75 wt% acetone, and 4.25 wt% water. The solution was added to a round bottom flask. The flask was rotated at approximately 150 rpm in a 40°C water bath under a reduced pressure of about 0.1 atm. The resulting solid dispersion was removed from the flask as fine granules and used without further processing.

Example 9 was prepared by spraying a coating solution comprising 2.5 wt% Drug 1, 7.5 wt% HPMCAS-MF, and 90 wt% solvent (5 wt% water in acetone) onto Nu-Core beads (45/60 mesh) to produce a coating of an amorphous solid dispersion of the drug and polymer on the surface of the beads. An analysis showed that the coated beads contained 3.9 wt% Drug 1.

The drug, polymer and solvents for Examples 8 and 9 are shown in Table 2.

Table 2

Ex. No.	Drug	Drug Mass	Polymer	Polymer Mass	Drug Conc. in the Dispersion (wt%)	Solv	Solv Mass
8	1	1.875g	HPMCAS-MF (rotoevaporated)	1.875g	50	5 wt% water in acetone	21.25g
9	1	20g	HPMCAS-MF (coated beads)	60g	25	5 wt% water in acetone	720g

## CONTROLS 1-2

Comparative compositions Control 1 and Control 2 were simply 3.6 mg of crystalline Drug 1 and 3.6 mg of the amorphous form of Drug 1 respectively.

## Example 10

*In vitro* dissolution tests were performed to evaluate the performance of the amorphous dispersions of Examples 1-9 relative to the performance of Controls 1 and 2. The dissolution performance of the SDD of Example 1 was evaluated in an *in vitro* dissolution test using a microcentrifuge method. In this test, 14.4 mg of the SDD of Example 1 was added to a microcentrifuge tube. The tube was placed in a 37°C sonicating bath, and 1.8 mL phosphate buffered saline (PBS) at pH 6.5 and 290 mOsm/kg was added. The samples were quickly mixed using a vortex mixer for about 60 seconds. The samples were centrifuged at 13,000 G at 37°C for 1 minute. The resulting supernatant solution was then sampled and diluted 1:6 (by volume) with methanol and then analyzed by high-performance liquid chromatography (HPLC). The contents of the tubes were mixed on the vortex mixer and allowed to stand undisturbed at 37°C until the next sample was taken. Samples were collected at 4, 10, 20, 40, 90, and 1200 minutes.

The performance of Example Nos. 2-8 was likewise evaluated in *in vitro* dissolution tests using the same microcentrifuge method described above. The

dosage for each of these tests was 2000  $\mu\text{g}/\text{ml}$ . The results of the dissolution tests are shown in Table 3.

The performance of the amorphous dispersions of Example 9 were tested using the same microcentrifuge method, except that 2.5 grams of the coated beads were added to 50 mL of PBS solution (resulting in a dosage of 2000  $\mu\text{g}/\text{mL}$ ).

For Controls 1 and 2, *in vitro* dissolution tests were also performed using the same microcentrifuge method except that 3.6 mg of either crystalline or amorphous Drug 1 was used.

Table 3

Example	Time (mins)	Drug Concentration ( $\mu\text{g/mL}$ )	AUC (min* $\mu\text{g/mL}$ )
1	0	0	0
	4	635	1,300
	10	644	5,100
	20	711	11,900
	40	769	26,700
	90	844	67,000
	1200	1290	1,251,400
2	0	0	0
	4	601	1,200
	10	625	4,900
	20	653	11,300
	40	624	24,000
	90	693	57,000
	1200	548	745,700
3	0	0	0
	3	544	1,100
	10	558	4,400
	20	558	9,980
	40	552	21,100
	90	565	49,000
4	0	0	0
	3	526	1,100
	10	637	4,500
	20	649	11,000
	40	651	24,000
	90	688	57,400
5	0	0	0
	3	2066	4,100
	10	2035	16,400
	20	2075	37,000
	40	1965	77,400
	90	1845	173,600
6	0	0	0
	3	2040	4,100
	10	1777	15,500
	20	1704	32,900
	40	1483	64,800
	90	427	112,600
7	0	0	0
	3	1036	2,100
	10	1277	9,000
	20	1246	21,600
	40	1217	46,300
	90	503	89,300
	1200	350	562,700

Example	Time (mins)	Drug Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
8	0	0	0
	4	134	270
	10	197	1,300
	20	248	3,500
	40	308	9,100
	90	378	26,200
	1200	591	564,000
9	0	0	0
	4	412	820
	10	491	3,500
	20	523	8,600
	40	561	19,400
	90	617	48,900
	180	752	110,500
	1200	967	928,000
Control 1	0	0	0
	4	130	260
	10	149	1,100
	20	139	2,500
	40	149	5,400
	90	147	12,800
	1200	125	163,800
Control 2	0	0	0
	4	586	1,200
	10	473	4,400
	20	220	7,800
	40	182	11,700
	90	167	20,600
	180	158	35,200
	1200	203	225,900

The results of the *in vitro* dissolution tests  
 are summarized in Table 4, which shows the maximum  
 concentration of Drug 1 in solution during the first 90  
 minutes of the test ( $C_{\text{max},90}$ ), the area under the aqueous  
 concentration versus time curve after 90 minutes ( $\text{AUC}_{90}$ ),  
 and the concentration at 1200 minutes ( $C_{1200}$ ).

Table 4

Example	Dosage ( $\mu\text{g/mL}$ )	$C_{\text{max},90}$ ( $\mu\text{g/mL}$ )	$\text{AUC}_{90}$ ( $\text{min} \cdot \mu\text{g/mL}$ )	$C_{1200}$ ( $\mu\text{g/mL}$ )
1	2000	844	67,000	1290
2	2000	693	57,000	548
3	2000	565	49,000	397
4	2000	688	57,400	409
5	2000	2075	172,600	255
6	2000	2040	112,600	257
7	2000	1277	89,300	350
8	2000	378	26,200	591
9	2000	617	48,900	967
Control 1	2000	149	12,800	125
Control 2	2000	586	20,600	203

The results, summarized in Table 4, show that the performance of the SDDs of Examples 1-9 was much better than that of the crystalline drug alone (Control 1), with  $C_{\text{max},90}$  values ranging from 2.5- to 13.9-fold that of the crystalline drug, Control 1, and  $\text{AUC}_{90}$  values ranging from 2- to 13.4-fold that of the crystalline drug, Control 1. With respect to the amorphous drug alone, the dispersions of Examples 1-9 demonstrated an  $\text{AUC}_{90}$  that was 1.27- to 8.4-fold that of the amorphous drug alone, Control 2.

#### Example 11

This example shows improved *in vivo* performance of an amorphous dispersion of Drug 1 and concentration-enhancing polymer compared with the crystalline form of Drug 1. For Example 11, an SDD was prepared following the procedure described in Example 1. The SDD was then formulated as an oral powder for constitution (OPC) by suspending 1.2 gm of the SDD in 100 ml of a 0.5 wt% solution of Polysorbate 80 in sterile water. This OPC, which contained 300 mg of active Drug 1, was taken orally by healthy human subjects (n=4). The dosing bottle was rinsed twice with 100 ml of sterile water and administered orally to the subjects. As a control

(Control 3), an OPC was formed using an equivalent quantity of the crystalline form of Drug 1. The results of these *in vivo* tests are shown in Table 5, giving the maximum concentration of drug achieved in the blood plasma, the time to reach this maximum concentration, and the blood plasma drug AUC from 0 to 24 hours.

Table 5

Ex. No.	Dose (mg)	C <sub>max</sub> (μg/mL)	Time to C <sub>max</sub> (hr)	AUC <sub>0-24</sub> (μg-hr/mL)
11	300	8.4±1.1	2.5±0.6	46±7.6
Control 3	300	1.3±0.3	2.3±1.3	7.4±3.3

As shown in Table 5, the OPC of Example 11 showed improved performance compared with the OPC of Control 3, thus demonstrating the advantage of using an amorphous dispersion of a GPI and concentration-enhancing polymer. Not only was the blood plasma C<sub>max</sub> for Example 11 6.5-fold the blood plasma C<sub>max</sub> for Control 3, but the blood plasma AUC<sub>0-24</sub> for Example 11 was 6.21-fold that of Control 3.

#### Examples 12-17

These examples demonstrate the utility of the GPI amorphous dispersions of the present invention with another GPI, 5-chloro-1H-indole-2-carboxylic acid [1S-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl] amide ("Drug 2"), which has a solubility in water of 14.6 μg/mL. For Example 12, a solution containing 0.5 wt% Drug 2 and 0.5 wt% HPMCAS-LF in acetone was prepared. This solution was pumped into a "mini" spray-dryer apparatus via a syringe pump at a rate of 1.3 mL/min. The polymer solution was atomized through a spray nozzle using a heated stream of nitrogen (100°C). The resulting solid SDD containing 50 wt% Drug 2 was collected on filter paper at a yield of about 80%.

Examples 13-17 were prepared using the same method used to prepare Example 12, but with different polymers and in some cases different solvents. The variations are noted in Table 6.

Table 6

Ex. No.	Drug	Drug Mass	Polymer*	Polymer Mass	Drug Conc. in the Dispersion (wt%)	Solv	Solv Mass	Spray Dryer
12	2	25mg	HPMCAS-LF	25mg	50	acetone	5g	mini
13	2	15mg	HPMCAS-LF	45mg	25	methanol	10g	mini
14	2	15mg	HPMCP-55	45mg	25	methanol	10g	mini
15	2	15mg	PVP	45mg	25	10 wt% water in methanol	11g	mini
16	2	150mg	CAP	150mg	50	50 wt% water in acetone	33.4 g	mini
17	2	150mg	CAT	150mg	50	50 wt% water in acetone	33.4 g	mini

\* Polymer designations: HPMCAS = hydroxypropyl methyl cellulose acetate succinate, PVP = polyvinylpyrrolidone, CAP = cellulose acetate phthalate, CAT = cellulose acetate trimellitate, HPMCP = hydroxypropyl methyl cellulose phthalate.

## CONTROLS 4-5

Comparative compositions Control 4 and Control 5 were simply 1.8 mg of crystalline Drug 2 and 1.8 mg of amorphous Drug 2, respectively.

## Example 18

*In vitro* dissolution tests were performed to evaluate the performance of the amorphous dispersions of Examples 12-17 relative to the performance of Controls 4 and 5. The SDD of Example 12 was evaluated in an *in vitro* dissolution test using a microcentrifuge method. In this test, 3600  $\mu$ g of the SDD of Example 12 was added to a microcentrifuge tube. The tube was placed in a 37°C sonicating bath, and 1.8 mL of model fasted duodenal solution (MFDS), comprising phosphate buffered saline



with 14.7 mM sodium taurocholic acid and 2.8 mM of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine, pH 6.5, 290 mOsm/kg, was added. This resulted in a dose of Drug 2 of 1000  $\mu\text{g}/\text{mL}$ . The samples were quickly mixed using a vortex mixer for about 60 seconds. The samples were centrifuged at 13,000 G at 37°C for 1 minute. The resulting supernatant solution was then sampled and diluted 1:6 (by volume) with methanol and then analyzed by high-performance liquid chromatography (HPLC). The contents of the tubes were mixed on the vortex mixer and allowed to stand undisturbed at 37°C until the next sample was taken. Samples were collected at 4, 10, 20, 40, 90, and 1200 minutes.

For Controls 4 and 5, *in vitro* dissolution tests were performed using the procedures described above except that 1.8 mg of crystalline and amorphous Drug 2 was used, respectively.

Results of the dissolution tests are presented in Table 7.

Table 7

Example	Time (mins)	Drug Concentration ( $\mu\text{g}/\text{mL}$ )	AUC ( $\text{min} \cdot \mu\text{g}/\text{mL}$ )
12	0	0	0
	4	850	1,700
	10	866	6,900
	20	895	15,700
	40	908	33,700
	90	923	79,500
	1200	933	1,109,500
13	0	0	0
	4	958	1,900
	10	993	7,800
	20	997	17,700
	40	975	37,400
	90	992	86,600
	1200	961	1,170,500
14	0	0	0
	4	860	1,700
	10	822	6,800
	20	804	14,900
	40	805	31,000
	90	778	70,600
	1200	732	908,600

Example	Time (mins)	Drug Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
15	0	0	0
	4	467	930
	10	498	3,800
	20	505	8,800
	40	507	19,000
	90	495	44,000
	1200	545	621,200
16	0	0	0
	4	696	1,400
	10	708	5,600
	20	708	12,700
	40	695	26,700
	90	701	61,600
	1200	735	858,600
17	0	0	0
	4	768	1,500
	10	766	6,100
	20	746	13,700
	40	730	28,500
	90	744	65,300
	1200	722	878,900
Control 4	0	0	0
	4	45	90
	10	44	357
	20	53	842
	40	72	2,100
	90	82	5,900
	1200	102	108,100
Control 5	0	0	0
	4	151	302
	10	203	1,400
	20	225	3,500
	40	238	8,100
	90	268	20,800
	1200	370	374,900

The results of these tests are summarized in Table 8, which shows the maximum concentration of Drug 2 in solution during the first 90 minutes of the test ( $C_{\text{max},90}$ ), the aqueous area under the curve after 90 minutes ( $\text{AUC}_{90}$ ), and the concentration at 1200 minutes ( $C_{1200}$ ).

Table 8

Example	Dosage ( $\mu\text{g/mL}$ )	$C_{\text{max}, 90}$ ( $\mu\text{g/mL}$ )	$\text{AUC}_{90}$ ( $\text{min} \cdot \mu\text{g/mL}$ )	$C_{1200}$ ( $\mu\text{g/mL}$ )
12	1000	923	79,500	933
13	1000	997	86,600	961
14	1000	860	70,600	732
15	1000	507	44,000	545
16	1000	708	61,600	735
17	1000	768	65,300	722
Control 4	1000	82	5,900	102
Control 5	1000	268	20,800	370

In general, the dispersions of Examples 12-17 showed much better performance than the crystalline drug alone, with  $C_{\text{max}, 90}$  values ranging from 6.2- to 12.1-fold that of the crystalline drug, Control 4, and  $\text{AUC}_{90}$  values ranging from 7.5- to 14.7-fold that of the crystalline drug, Control 4. With respect to the amorphous drug alone, all of the dispersions of Examples 12-17 demonstrated a  $C_{\text{max}}$  and an  $\text{AUC}_{90}$  greater than that of the amorphous drug alone, with  $C_{\text{max}, 90}$  values ranging from 1.9- to 3.7-fold that of the amorphous drug, Control 5, and  $\text{AUC}_{90}$  values ranging from 2.1- to 4.2-fold that of the amorphous drug, Control 5.

#### Example 19

This example demonstrates that the compositions of this invention, when orally dosed to beagle dogs, give a high systemic compound exposure ( $C_{\text{max}}$  and  $\text{AUC}$ ). An amorphous solid dispersion of 50 wt% Drug 2 and 50 wt% polymer was made by first mixing Drug 2 in the solvent acetone together with HPMCAS-LF to form a solution. The solution comprised 2.5 wt% Drug 2, 2.5 wt% HPMCAS-LF, and 95 wt% acetone. This solution was then spray-dried by directing an atomizing spray using a two-fluid external-mix spray nozzle at 2.2 bar at a feed rate of 200 g/min into the stainless-steel chamber of a Niro PSD-1 spray-dryer, maintained at a temperature of 180°C at the inlet and 68°C at the outlet.

The resulting amorphous solid SDD was collected via a cyclone and then dried in a Gruenberg solvent tray-dryer by spreading the spray-dried particles onto polyethylene-lined trays to a depth of not more than 1 cm and then drying them at 40°C for at least 8 hours.

The SDD was dosed as an oral powder for constitution (OPC) by suspending 200 mg of the SDD in approximately 20 ml of a 2 wt% solution of Polysorbate 80 in sterile water. This OPC, containing 100 mg of active Drug 2 was administered orally to beagle dogs using an oral gavage tube. As a control (Control 6), a similar OPC was formed using the crystalline form of the drug. Relative bioavailability was calculated by dividing the AUC in the blood of subjects receiving the test dose by the AUC in the blood of subjects receiving the control dose (Control 6).

Dogs that had fasted overnight were dosed with suspensions containing 100 mg of Drug 2, along with 20 mL of water. Blood was collected from the jugular vein of the dogs before dosing and at various time points after dosing. To 100  $\mu$ L of each plasma sample, 5 mL of methyl-tert-butyl ether (MTBE) and 1 mL of 500 mM sodium carbonate buffer (pH 9) were added; the sample was vortexed for 1 minute and then centrifuged for 5 minutes. The aqueous portion of the sample was frozen in a dry-ice/acetone bath, and the MTBE layer was decanted and evaporated in a vortex evaporator. Dried samples were reconstituted in 100  $\mu$ L of mobile phase (33% acetonitrile and 67% of 0.1% formic acid in water). Analysis was carried out by HPLC. The results of these tests are shown in Table 9, where  $C_{\max}$  is the maximum concentration in the blood plasma,  $AUC_{0-24}$  is the area under the drug concentration in the blood curve in the first 24 hours, and Relative Bioavailability is the AUC in the blood of subjects receiving the test dose divided by the AUC of subjects receiving the Control 6.

Table 9

Example No.	C <sub>max</sub> , ( $\mu\text{g/mL}$ )	AUC <sub>0-24</sub> ( $\mu\text{g-hr/mL}$ )	Relative Bioavailability
19	9.8 $\pm$ 4.6	38 $\pm$ 6	6.2
Control 6	1.6 $\pm$ 0.7	6.1 $\pm$ 4.0	1

The results show the superior performance of the amorphous GPI and polymer dispersion of Example 19 relative to the crystalline GPI, Control 6, providing a C<sub>max</sub> value that was 6.1-fold that of the control and a relative bioavailability of 6.2 relative to the control.

## Examples 20-25

Examples 20-25 demonstrate the utility of the GPI amorphous dispersions of the present invention with another GPI, 5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-methoxy-methylcarbamoylmethyl)-2-phenylethyl]-amide ("Drug 3"), which has a solubility in water of 1  $\mu\text{g/mL}$  and a solubility in MFD solution of 17  $\mu\text{g/mL}$ . To prepare Example 20, a solution containing 0.5 wt% of Drug 3 and 0.5 wt% HPMCAS-MF in acetone was prepared. This solution was pumped into a "mini" spray-dryer apparatus via a syringe pump at a rate of 1.3 mL/min. The polymer solution was atomized through a spray nozzle using a heated stream of nitrogen (100°C). The resulting solid SDD containing 50 wt% Drug 3 was collected on a filter paper at a yield of about 62%.

Examples 21-25 were prepared using the same method used to prepare Example 20, but with different polymers and in some cases different solvents. The variations are note in Table 10.

Table 10

Ex. No.	Drug	Drug Mass	Polymer*	Polymer Mass	Drug Conc. in the Dispersion (wt%)	Solv	Solv Mass	Spray Dryer
20	3	52 mg	HPMCAS-MF	52 mg	50.0	acetone	12 g	mini
21	3	50.5 mg	PVP	50.4 mg	50.0	acetone methanol	12 g 0.24 g	mini
22	3	49.7 mg	HPMCP	49.9 mg	49.9	acetone	12 g	mini
23	3	50.1 mg	CAP	50.3 mg	49.9	acetone	12 g	mini
24	3	50.9 mg	HPC	51.8 mg	49.6	acetone	12 g	mini
25	3	50 mg	PVAP	50 mg	50.0	acetone	12 g	mini

\* Polymer designations: HPMCAS = hydroxypropyl methyl cellulose acetate succinate, HPMC = hydroxypropyl methyl cellulose, PVP = polyvinylpyrrolidone, CAP = cellulose acetate phthalate, HPC = hydroxypropyl cellulose, PVAP = polyvinyl acetate phthalate, HPMCP = hydroxypropyl methyl cellulose phthalate.

## Control 8

Comparative composition Control 8 consisted of 5 mg of the crystalline form of Drug 3 alone.

## Example 26

In vitro dissolution tests were performed to evaluate the performance of the amorphous dispersions of Examples 20-25 relative to the performance of Control 8. The SDD of Example 20 was evaluated in an in vitro dissolution test using a syringe/filter method. In this test, 10 mg of the SDD of Example 20 was added to 10 mL of MFD solution, comprising phosphate buffered saline with 14.7 mM sodium taurocholic acid and 2.8 mM of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine, pH 6.5, 290 mOsm/kg. The drug solution was added to a 10 mL polypropylene syringe fitted with a Titan PVDF 0.45  $\mu$ m filter. The syringe was attached to a vertical rotating wheel in a 37°C constant temperature chamber. At each sampling time, 13 drops were expelled from the syringe through the filter. The filtrate was then diluted 1:1 (by volume) with methanol and analyzed by high-performance liquid chromatography (HPLC). Between sampling times,

the test solution was mixed as the syringe was rotated on the wheel at 37°C. Samples were collected at 0.5, 5, 30, 60, 180, and 1200 minutes.

5        *In vitro* dissolution tests for Examples 21-25 were performed using the same procedure described above for Example 20.

For Control 8, an *in vitro* dissolution test was performed using the procedure described above except that 5 mg of crystalline Drug 3 was used.

10        The concentrations of drug obtained in these samples are shown in Table 11 below.

Table 11

Example	Time (mins)	Drug Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
20	0	0	0
	0.5	25	6
	5	62	202
	30	112	2,400
	60	115	5,800
	180	120	19,900
	1200	14	88,200
21	0	0	0
	0.5	8	2
	5	33	94
	30	121	2,000
	60	128	5,800
	180	114	20,300
	1200	13	85,000
22	0	0	0
	0.5	12	3
	5	48	138
	30	112	2,100
	60	128	5,700
	180	106	19,800
	1200	13	80,500
23	0	0	0
	0.5	14	4
	5	46	139
	30	106	2,000
	60	121	5,400
	180	127	20,300
	1200	13	91,700
24	0	0	0
	0.5	24	6
	5	62	200
	30	94	2,200
	60	95	5,000
	180	91	16,100
	1200	11	68,200
25	0	0	0
	0.5	6	2
	5	34	92
	30	89	1,600
	60	104	4,500
	180	17	11,800
	1200	9	25,000
Control 8	0	0	0
	0.5	3	1
	5	8	26
	30	10	251
	60	10	551
	180	9	1,700
	1200	8	10,400



The results of these test are summarized in Table 12, which shows the maximum concentration of Drug 3 in solution after 180 minutes ( $C_{\max 180}$ ), the aqueous area under the curve after 180 minutes ( $AUC_{180}$ ), and the concentration at 1200 minutes ( $C_{1200}$ ).

Table 12

Example	Dosage ( $\mu\text{g/mL}$ )	$C_{\max 180}$ ( $\mu\text{g/mL}$ )	$AUC_{180}$ ( $\text{min} \cdot \mu\text{g/mL}$ )	$C_{1200}$ ( $\mu\text{g/mL}$ )
20	500	120	19,900	14
21	500	133	20,300	13
22	500	127	19,800	13
23	500	127	20,300	13
24	500	97	16,100	11
25	500	104	11,800	9
Control 8	500	10	1,700	8

The results show that the performance of the SDD of Examples 20-25 was much better than that of the crystalline drug alone, with  $C_{\max 180}$  values 9.7- to 13.3-fold that of Control 8, and  $AUC_{180}$  values 6.9- to 11.9-fold that of Control 8.

## Examples 27-29

These examples disclose simple physical mixtures of a GPI and a concentration-enhancing polymer. Mixtures of Drug 1 and HPMCAS-MF were formed by dry mixing amorphous Drug 1 with HPMCAS-MF. For Example 27, the composition comprised 3.6 mg (75 wt%) Drug 1 and 1.2 mg (25 wt%) HPMCAS-MF; for Example 28, the composition comprised 3.6 mg (50 wt%) Drug 1 and 3.6 mg (50 wt%) HPMCAS-MF; for Example 29, the composition comprised 3.6 mg (25 wt%) Drug 1 and 10.8 mg (75 wt%) HPMCAS-MF.

These compositions were evaluated in *in vitro* dissolution tests using the procedures outlined in Example 10. The quantities of drug and polymer noted above were each added to a microcentrifuge tube, to which

was added 1.8 ml of PBS solution. The tube was vortexed immediately after adding the PBS solution. The results of these dissolution tests are given in Table 13, and summarized in Table 14.

5

Table 13

Example	Time (mins)	Drug 1 Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
27	0	0	0
	4	714	1,400
75 wt% Drug 1/	10	737	5,800
25 wt% HPMCAS-MF	20	696	12,900
	40	690	26,800
	90	729	62,300
	180	684	125,800
	1200	440	696,600
28	0	0	0
	4	377	755
50 wt% Drug 1/	10	370	3,000
50 wt% HPMCAS-MF	20	836	9,000
	40	846	25,800
	90	898	69,500
	180	918	151,200
	1200	627	932,700
29	0	0	0
	4	999	2,000
25 wt% Drug 1/	10	1030	8,100
75 wt% HPMCAS-MF	20	1065	18,600
	40	1133	40,600
	90	1185	98,500
	180	1304	210,500
	1200	1379	1,579,500

30

Table 14

Example	Dosage ( $\mu\text{g/mL}$ )	$C_{\text{max}, 90}$ ( $\mu\text{g/mL}$ )	$AUC_{90}$ ( $\text{min} \cdot \mu\text{g/mL}$ )	$C_{1200}$ ( $\mu\text{g/mL}$ )
27	2000	729	62,300	440
28	2000	898	69,500	627
29	2000	1185	98,500	1379
Control 2	2000	586	20,600	203

40

These simple physical mixtures of amorphous Drug 1 and HPMCAS-MF showed much better performance than the amorphous drug alone (Control 2, shown in Table 14 for comparison), with  $C_{\text{max}, 90}$  values that were 1.24- to

2.0-fold that of Control 2, and AUC<sub>90</sub> values that were 3.0- to 4.8-fold that of Control 2.

#### Example 30

5           This example demonstrates another simple physical mixture of amorphous GPI and polymer. A coating solution comprising 7.5 wt% HPMCAS-MF dissolved in 92.5 wt% solvent (5 wt% water in acetone) was prepared and spray-coated onto Nu-Core Beads (45/60 mesh),  
10           producing a thin coating of the polymer on the surface of the beads resulting in beads containing 12.2 wt% HPMCAS-MF. Samples of these beads (2.4 gm) were then mixed with 100 mg of amorphous Drug 1 (resulting in a drug:polymer ratio of 1:3 or 25 wt% Drug 1) and evaluated  
15           in an *in vitro* dissolution test using the procedures outlined in Example 10. The results of the dissolution test are presented in Table 15.

Table 15

20

Example	Time (mins)	Drug 1 Concentration ( $\mu\text{gA/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
30	0	0	0
	4	797	1,600
	10	1047	7,100
	20	1292	18,800
	40	1523	47,000
	90	1653	126,400
	180	1724	278,300
	1200	1885	2,113,600

25

          The physical mixture of HPMCAS-MF coated beads with amorphous Drug 1 showed improved performance over crystalline Drug 1 alone, with a C<sub>max,90</sub> value that is  
30           11-fold that of crystalline Drug 1 (Control 1) and an AUC<sub>90</sub> value that is 9.9-fold that of Control 1.

#### Example 31

          A composition was formed by blending 50 wt% of  
35           the SDD of Example 2 (containing 50 wt% Drug 1 and 50 wt%

HPMCAS-MF) with 50 wt% HPMCAS-MF. This composition was evaluated in a dissolution test as described in Example 10. The results of this test are presented in Table 16, and show that the blend of the SDD with polymer performs well, with a  $C_{\max, 90}$  value that is 6.6-fold that of the crystalline drug alone (Control 1) and an  $AUC_{90}$  value that is 6.2-fold that of Control 1.

Table 16

Example	Time (mins)	Drug 1 Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
31	0	0	0
	4	766	1,500
	10	840	6,400
	20	874	14,900
	40	884	32,500
	90	979	79,100
	1200	1133	1,251,000

## Examples 32-35

An amorphous solid dispersion of 50 wt% Drug 1 and 50 wt% polymer was made by first mixing Drug 1 in a solvent together with HPMCAS-MF to form a solution. The solution comprised 7.5 wt% Drug 1, 7.5 wt% HPMCAS, 80.75 wt% acetone and 4.25 wt% water. This solution was then spray-dried by directing an atomizing spray using a two-fluid external-mix spray nozzle at 2.7 bar (37 psig) at a feed rate of 175 g/min into the stainless-steel chamber of a Niro spray-dryer, maintained at a temperature of 175°C at the inlet and 70°C at the outlet.

The resulting amorphous solid spray-dried dispersion (SDD) was collected via a cyclone and then dried in a Gruenberg solvent tray-dryer by spreading the spray-dried particles onto polyethylene-lined trays to a depth of not more than 1 cm and then drying them at 40°C for 16 hours.

The SDD above was incorporated into tablets containing 25, 50, 100, and 200 mg. Tablets with a dose of 25 mg (Example 32) consisted of 7.14 wt% SDD, 40.0 wt%

HPMCAS-MF, 49.11 wt% microcrystalline cellulose (Avicel® PH 102), 3.0 wt% croscarmellose sodium (Ac-Di-Sol®), and 0.75 wt% magnesium stearate. Tablets with a dose of 50 mg (Example 33) consisted of 14.29 wt% SDD, 40.0 wt% HPMCAS-MF, 41.96 wt% Avicel® PH 102, 3.0 wt% Ac-Di-Sol®, and 0.75 wt% magnesium stearate. Tablets with a dose of 100 mg (Example 34) consisted of 28.57 wt% SDD, 30.0 wt% HPMCAS-MF, 37.68 wt% Avicel® PH 102, 3.0 wt% Ac-Di-Sol®, and 0.75 wt% magnesium stearate. Tablets with a dose of 200 mg (Example 35) consisted of 57.14 wt% SDD, 39.11 wt% Avicel® PH 102, 3.0 wt% Ac-Di-Sol®, and 0.75 wt% magnesium stearate. In each case, the targeted tablet weight was 700 mg.

To form the tablets, the SDD was first granulated (roller compacted) on a Freund TF-mini roller compactor using an auger speed of 30 rpm, a roller speed of 4 rpm, and a roller pressure of 30 Kg<sub>f</sub>/cm<sup>2</sup>. The resulting compacted material was then reduced using a mini-Comil at a power setting of 4, with sieve 039R. The milled SDD was then blended in a V-blender with the HPMCAS-MF, Avicel®, and Ac-Di-Sol® for 20 minutes using the proportions noted above. Next, a portion of the magnesium stearate (about 20 wt% of the total magnesium stearate used) was added and the material was blended for 5 minutes. The blend was then granulated again using an auger speed of 20 rpm, a roller speed of 4 rpm, and a roller pressure of 30 Kg<sub>f</sub>/cm<sup>2</sup>. The resulting compacted material was then reduced using a Comill with a power setting of 3 and a sieve size of 032R. The remaining magnesium stearate was then added, and the material was blended for 5 minutes in a V-blender. This material was then formed into tablets using 0.3437 x 0.6875-inch oval tooling on a Kilian T-100 tablet press with precompression of 1 to 2 kN and a compression force of 10 kN.

To test *in vitro* drug dissolution, one of each of the tablets was each placed in 200 mL of a gastric

buffer solution (0.1 N HCl at pH 1.2) for 30 minutes at 37°C and stirred, after which 50 mL of a pH 13 buffer solution was added to produce a final pH of 7.5 and a final volume of 250 mL. The drug concentration was determined over time by periodically withdrawing samples, centrifuging the samples to remove any undissolved drug, diluting the supernatant in methanol, analyzing the samples by HPLC, and calculating drug concentrations. The concentrations of drug obtained in *in vitro* dissolution tests are shown in Table 17 below.

Table 17

Example	Time (mins)	Drug 1 Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
32 25 mg	0	0	0
	5	6	16
	15	13	110
	20	15	178
	35	25	478
	45	30	755
	60	36	1,300
	75	43	1,800
	90	50	2,500
	120	58	4,200
	180	65	7,900
	1200	96	90,200
33 50 mg	0	0	0
	5	9	24
	15	19	166
	20	23	271
	35	42	755
	45	61	1,300
	60	82	2,300
	75	99	3,700
	90	111	5,300
	120	130	8,900
	180	152	17,400
	1200	202	197,800

Example	Time (mins)	Drug 1 Concentration ( $\mu\text{g/mL}$ )	AUC ( $\text{min} \cdot \mu\text{g/mL}$ )
34 100 mg	0	0	0
	5	20	49
	15	43	361
	20	50	594
	35	112	1,800
	45	150	3,100
	60	186	5,700
	75	199	8,500
	90	213	11,600
	120	236	18,300
	180	260	33,200
	1200	381	360,300
35 200 mg	0	0	0
	5	26	64
	15	64	514
	20	81	878
	35	168	2,800
	45	424	5,700
	60	470	12,400
	75	479	19,500
	90	502	26,900
	120	518	42,200
	180	522	73,400
	1200	298	491,000

5

The data demonstrate that approximately all of the drug had been released by 1200 minutes.

10

#### Example 36

An amorphous solid dispersion of 67 wt% Drug 3 and 33 wt% polymer was made by first mixing Drug 3 in the solvent acetone together with HPMCAS-MF to form a solution. The solution comprised 3.33 wt% Drug 3, 1.67 wt% HPMCAS-MF, and 95 wt% acetone. This solution was then spray-dried by directing an atomizing spray using a two-fluid external-mix spray nozzle at 0.6 bar at a feed rate of 75 g/min into the stainless-steel chamber of a Niro PSD-1 spray-dryer, maintained at a temperature of 120°C at the inlet and 76°C at the outlet.

20

The resulting amorphous solid spray-dried dispersion (SDD) was collected via a cyclone and then dried in a Gruenberg solvent tray-dryer by spreading the spray-dried particles onto polyethylene-lined trays to a

depth of not more than 1 cm and then drying them at 40°C for at least 8 hours.

#### Example 37

5 Capsules containing a total mass of 500 mg were prepared using the SDD of Drug 3 from Example 36. Each capsule contained 60 wt% of the SDD, 15 wt% Fast Flo lactose, 15 wt% Avicel PH-102, 7 wt% Explotab, 2 wt% sodium lauryl sulfate, and 1 wt% magnesium stearate,  
10 resulting in capsules containing 200 mg of Drug 3.

#### Example 38

Tablets with a total mass of 600 mg were prepared containing 50 wt% SDD from Example 36, 32 wt%  
15 Avicel PH-102, 11 wt% Fast Flo lactose, 5 wt% Explotab, 1 wt% sodium lauryl sulfate, and 1 wt% magnesium stearate, resulting in tablets containing 200 mg of Drug 3.

#### Examples 39-40

20 Capsules with a total mass of 600 mg were prepared, each capsule containing 50 wt% SDD from Example 36, 32 wt% Avicel PH-102, 11 wt% Fast Flo lactose, 5 wt% Explotab, 1 wt% sodium lauryl sulfate, and  
25 1 wt% magnesium stearate (Example 39), resulting in capsules containing 200 mg of Drug 3. Example 40 was prepared by coating the capsules of Example 38 with cellulose acetate phthalate.

#### Example 41

30 The dosage forms of Examples 37 to 40 were tested in *in vivo* tests. Beagle dogs that had fasted overnight were dosed with capsules and tablets from Examples 37 to 40, along with 50 mL of water. Blood was  
35 collected from the jugular vein of the dogs before dosing and at various time points after dosing. To 100  $\mu$ L of each plasma sample, 5 mL of methyl-tert-butyl ether



(MTBE) and 1 mL of 500 mM sodium carbonate buffer (pH 9) were added; the sample was vortexed for 1 minute and then centrifuged for 5 minutes. The aqueous portion of the sample was frozen in a dry-ice/acetone bath, and the MTBE layer was decanted and evaporated in a vortex evaporator. Dried samples were reconstituted in 100  $\mu$ L of mobile phase (33% acetonitrile and 67% of 0.1% formic acid in water). Analysis was carried out by HPLC.

As a control (Control 9), an OPC was formed using the crystalline form of Drug 3 as follows. An aqueous suspension of 200 mg of crystalline drug was prepared in 2 wt% Polysorbate 80 in water. Oral administration of the aqueous drug suspensions was facilitated using an oral gavage equipped with a polyethylene tube insert. The polyethylene tube insert was used to accurately deliver the desired volume of dose by displacement, without the need for additional volume of water to rinse the tube.

The results of these tests are shown in Table 18, where  $C_{\max}$  is the maximum concentration of Drug 3 in the blood plasma,  $AUC_{0-24}$  is the area under the curve in the first 24 hours, and Relative Bioavailability is the AUC in the blood of the test dose divided by the AUC in the blood of the reference dose (Control 9). The results show that the relative bioavailabilities obtained with the dosage forms of the present invention are 2.8 to 6.2 relative to Control 9. Furthermore, the  $C_{\max}$  of the dosage forms of the present invention were 2.6-fold to 4.7-fold that of Control 9.

Table 18

Example	Formulation	Dose (mg)	C <sub>max</sub> , (μg/mL)	AUC <sub>0-24</sub> (μg-hr/mL)	Relative Bioavailability
5 Control 9	Crystalline suspension	200	1.03 ± 0.57	6.48 ± 3.60	—
37	Capsule	200	3.81 ± 2.39	31.75 ± 18.61	6.0
38	Tablet	200	2.84 ± 2.04	26.09 ± 21.43	4.2
39	Capsule	200	4.86 ± 2.30	40.55 ± 20.74	6.2
10 40	CAP coated Capsule	200	2.67 ± 2.45	18.08 ± 12.21	2.8

## Example 42

This example illustrates a method for making a  
 15 tablet dosage form of the present invention containing an  
 amorphous dispersion of Drug 1. An amorphous solid  
 dispersion of Drug 1 and HPMCAS was made by mixing Drug 1  
 in a solvent together with HPMCAS to form a solution, and  
 then spray-drying the solution. The solution comprised  
 20 7.5 wt% Drug 1, 7.5 wt% HPMCAS-MF, 4.25 wt% water, and  
 80.75 wt% acetone. The solution was then spray-dried by  
 directing an atomizing spray using a two-fluid external-  
 mix spray nozzle at 2.7 bar at a feed rate of 175 g/min  
 into the stainless steel chamber of a Niro spray-dryer,  
 25 maintained at a temperature of 140°C at the inlet and 50°C  
 at the outlet. The resulting SDD was collected via a  
 cyclone and then dried in a Gruenberg solvent tray-dryer  
 by spreading the spray-dried particles onto polyethylene-  
 lined trays to a depth of not more than 1 cm and then  
 30 drying them at 40°C for at least 8 hours. After drying,  
 the SDD contained 50 wt% Drug 1.

The tablets contained 50 wt% SDD, 25 wt%  
 anhydrous dibasic calcium phosphate, 12 wt% Avicel®  
 PH 200, 12.5 wt% crospovidone, and 0.5 wt% magnesium  
 35 stearate. The total batch weight was 190 g. First, the  
 ingredients, except for magnesium stearate, were added to  
 a Turbula blender and blended for 20 minutes. Next, half  
 of the magnesium stearate was added and blended for  
 5 minutes. The blend was then roller-compacted with a

Vector TF mini roller compactor using an auger speed of 30 rpm, a roller speed of 5 rpm, and a roller pressure of 35.2 Kgf/cm<sup>2</sup>. The resulting compacted material was then milled using a Quadro Comil 193AS mill at a power setting of 3, using impeller 2B-1607-005 and Screen 2B-075R03151173. The second half of the magnesium stearate was added next, and the material was blended for 5 minutes in a Turbula blender. This material was then formed into 800 mg tablets using 1/2-inch SRC tooling on a Manesty F press. An average tablet hardness of 19 Kp was obtained. Average disintegration time in deionized water (USP disintegration apparatus) was 2 minutes, 50 seconds.

#### Example 43

The tablets of Example 42 were coated in a LDGS 20 pan-coater using an 8 wt% aqueous solution of Opadry® II Clear. The following coating conditions were used: tablet bed weight, 900 g; pan speed, 20 rpm; outlet temperature, 40°C; solution flow, 8 g/min; atomization pressure, 20 psi; and air flow, 40 cfm. The coating weight gain was 3 wt%. The resulting average coated tablet hardness was 45 Kp. Average disintegration time in deionized water was 4 minutes, 57 seconds.

25

#### Example 44

This example illustrates another method for making a tablet dosage form of the present invention containing an amorphous dispersion of Drug 1. An amorphous solid dispersion of Drug 1 and HPMCAS was made by mixing Drug 1 in a solvent together with HPMCAS to form a solution, and then spray-drying the solution, as described in Example 42. The tablets contained 50 wt% of the SDD, 25 wt% anhydrous dibasic calcium phosphate, 12 wt% Avicel® PH 105 QS, 12.5 wt% crospovidone, and 0.5 wt% magnesium stearate. To form the tablets, the ingredients, except magnesium stearate, were first added

35

to a V-blender and blended for 20 minutes, followed by de-lumping using a 10-mesh screen. Next, half of the magnesium stearate was added and blended for 5 minutes. The blend was then roller compacted with a Vector TF mini roller compactor, fitted with "S"-type rolls, using an  
5     auger speed of 30 rpm, a roller speed of 4 rpm, and a roller pressure of 30 Kgf/cm<sup>2</sup>. The resulting compacted material was then milled using a Fitzpatrick M5A mill at a power setting of 350 rpm, with a sieve size of 16 mesh.  
10    The second half of the magnesium stearate was added next, and the material was blended for 5 minutes in a V-blender. This material was then formed into 800 mg tablets using 1/2-inch SRC tooling on a Killian T-100 (feeder frame speed 30 rpm, 30,000 tablets/hour), and  
15    compressed to a hardness of 25 Kp.

      The tablets above were coated in a Freund HCT-30 pan-coater using an aqueous solution of 3.5 wt% Opadry® II White and 0.5 wt% Opadry® II Clear. The following coating conditions were used: tablet bed  
20    weight, 1000 g; pan speed, 17 rpm; outlet temperature, 42°C; and solution flow, 6 g/min. Average disintegration time in deionized water was <5 minutes.

      The terms and expressions which have been employed in the foregoing specification are used therein  
25    as terms of description and not of limitation, and there is no intention, in the use of such terms and expressions, of excluding equivalents of the features shown and described or portions thereof, it being recognized that the scope of the invention is defined and  
30    limited only by the claims which follow.

## CLAIMS

1. A pharmaceutical composition comprising a  
glycogen phosphorylase inhibitor and a concentration-  
enhancing polymer wherein a portion of said glycogen  
phosphorylase inhibitor binds to a portion or all  
portions of the following residues of a glycogen  
phosphorylase enzyme:


10	parent secondary	
	<u>structure</u>	<u>residue number</u>
		13-23
	helix $\alpha$ 1	24-37
15	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80
	strand $\beta$ 1	81-86
		87-88
20	strand $\beta$ 2	89-92
		93
	helix $\alpha$ 3	94-102
		103
	helix $\alpha$ 4	104-115
25		116-117
	helix $\alpha$ 5	118-124
		125-128
	strand $\beta$ 3	129-131
		132-133
30	helix $\alpha$ 6	134-150
		151-152
	strand $\beta$ 4	153-160
		161

	strand $\beta$ 4b	162-163
		164-166
	strand $\beta$ 5	167-171
		172-173
5	strand $\beta$ 6	174-178
		179-190
	strand $\beta$ 7	191-192
		194, 197
	strand $\beta$ 8	198-209
10		210-211
	strand $\beta$ 9	212-216
	strand $\beta$ 10	219-226, 228-232
		233-236
	strand $\beta$ 11	237-239, 241, 243-247
15		248-260
	helix $\alpha$ 7	261-276
	strand $\beta$ 11b	277-281
	reverse turn	282-289
	helix $\alpha$ 8	290-304.

20

2. A pharmaceutical composition comprising a  
glycogen phosphorylase inhibitor and a concentration-  
enhancing polymer, said glycogen phosphorylase inhibitor  
being selected from the group consisting of Formula I,  
25 Formula II, Formula III and Formula IV;  
wherein Formula I is

30



35

Formula I

or the pharmaceutically acceptable salts or prodrugs thereof wherein the dotted line (---) is an optional bond wherein;

A is -C(H)=, -C((C<sub>1</sub>-C<sub>4</sub>)alkyl)= or -C(halo)= when  
5 the dotted line (---) is a bond, or A is methylene or -CH((C<sub>1</sub>-C<sub>4</sub>)alkyl)- when the dotted line (---) is not a bond;

R<sub>1</sub>, R<sub>10</sub> or R<sub>11</sub> are each independently H, halo, 4-, 6- or 7-nitro, cyano, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy,  
10 fluoromethyl, difluoromethyl or trifluoromethyl;

R<sub>2</sub> is H;

R<sub>3</sub> is H or (C<sub>1</sub>-C<sub>5</sub>)alkyl;

R<sub>4</sub> is H, methyl, ethyl, n-propyl, hydroxy(C<sub>1</sub>-C<sub>3</sub>)alkyl, (C<sub>1</sub>-C<sub>3</sub>)alkoxy(C<sub>1</sub>-C<sub>3</sub>)alkyl,  
15 phenyl(C<sub>1</sub>-C<sub>4</sub>)alkyl, phenylhydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, phenyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, thien-2- or -3-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl or fur-2- or -3-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl wherein said R<sub>4</sub> rings are mono-, di- or tri-substituted independently on carbon with H, halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl,  
20 (C<sub>1</sub>-C<sub>4</sub>)alkoxy, trifluoromethyl, hydroxy, amino or cyano; or

R<sub>4</sub> is pyrid-2-, -3- or -4-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, thiazol-2-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, imidazol-1-, -2-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrrol-2- or -3-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl,  
25 oxazol-2-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrazol-3-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, isoxazol-3-, -4-, -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, isothiazol-3-, -4-, -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyridazin-3- or -4-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrimidin-2-, -4-, -5- or -6-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrazin-2- or -3-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl or  
30 1,3,5-triazin-2-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, wherein said preceding R<sub>4</sub> heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino or hydroxy and said mono- or di-substituents are bonded to carbon;

35 R<sub>5</sub> is H, hydroxy, fluoro, (C<sub>1</sub>-C<sub>5</sub>)alkyl, (C<sub>1</sub>-C<sub>5</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkanoyl, amino(C<sub>1</sub>-C<sub>4</sub>)alkoxy, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkoxy,

carboxy(C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>5</sub>)alkoxy-carbonyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy, benzyloxycarbonyl(C<sub>1</sub>-C<sub>4</sub>)alkoxy, or carbonyloxy wherein said carbonyloxy is carbon-carbon linked with phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding R<sub>5</sub> rings are optionally monosubstituted with halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy, amino or trifluoromethyl and said mono-

substituents are bonded to carbon;

R<sub>7</sub> is H, fluoro or (C<sub>1</sub>-C<sub>5</sub>)alkyl; or  
R<sub>5</sub> and R<sub>7</sub> taken together are oxo;  
R<sub>6</sub> is carboxy or (C<sub>1</sub>-C<sub>8</sub>)alkoxycarbonyl, C(O)NR<sub>8</sub>R<sub>9</sub> or C(O)R<sub>12</sub> wherein

R<sub>8</sub> is H, (C<sub>1</sub>-C<sub>3</sub>)alkyl, hydroxy or (C<sub>1</sub>-C<sub>3</sub>)alkoxy; and

R<sub>9</sub> is H, (C<sub>1</sub>-C<sub>8</sub>)alkyl, hydroxy, (C<sub>1</sub>-C<sub>8</sub>)alkoxy, methylene-perfluorinated(C<sub>1</sub>-C<sub>8</sub>)alkyl, phenyl, pyridyl, thienyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl or 1,3,5-triazinyl wherein said preceding R<sub>9</sub> rings are carbon-nitrogen linked; or

R<sub>9</sub> is mono-, di- or tri-substituted (C<sub>1</sub>-C<sub>5</sub>)alkyl, wherein said substituents are independently H, hydroxy, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>5</sub>)alkylamino; or

R<sub>9</sub> is mono- or di-substituted (C<sub>1</sub>-C<sub>5</sub>)alkyl, wherein said substituents are independently phenyl, pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, pyridinyl, piperidinyl, morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl or 1,3,5-triazinyl

wherein the nonaromatic nitrogen-containing R<sub>9</sub> rings are optionally mono-substituted on nitrogen with (C<sub>1</sub>-C<sub>6</sub>)alkyl, benzyl, benzoyl or (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl and



wherein the R<sub>9</sub> rings are optionally mono-substituted on carbon with halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, hydroxy, amino, or mono-N- and di-N,N (C<sub>1</sub>-C<sub>5</sub>)alkylamino provided that no quaternized nitrogen is included and there are no  
5 nitrogen-oxygen, nitrogen-nitrogen or nitrogen-halo bonds;

R<sub>12</sub> is piperazin-1-yl, 4-(C<sub>1</sub>-C<sub>4</sub>)alkylpiperazin-1-yl, 4-formylpiperazin-1-yl, morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxo-thiomorpholino,  
10 thiazolidin-3-yl, 1-oxo-thiazolidin-3-yl, 1,1-dioxo-thiazolidin-3-yl, 2-(C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonylpyrrolidin-1-yl, oxazolidin-3-yl or 2(R)-hydroxymethylpyrrolidin-1-yl; or

R<sub>12</sub> is 3- and/or 4-mono- or di-substituted oxazetidin-2-yl, 2-, 4-, and/or 5- mono- or  
15 di-substituted oxazolidin-3-yl, 2-, 4-, and/or 5- mono- or di-substituted thiazolidin-3-yl, 2-, 4- and/or 5- mono- or di-substituted 1-oxothiazolidin-3-yl, 2-, 4-, and/or 5- mono- or di-substituted 1,1-dioxothiazolidin-3-yl, 3- and/or 4- mono- or di-substituted  
20 pyrrolidin-1-yl, 3-, 4- and/or 5-, mono-, di- or tri-substituted piperidin-1-yl, 3-, 4-, and/or 5- mono-, di-, or tri-substituted piperazin-1-yl, 3-substituted azetidin-1-yl, 4- and/or 5-, mono- or di-substituted 1,2-oxazinan-2-yl, 3- and/or 4- mono- or di-substituted  
25 pyrazolidin-1-yl, 4- and/or 5-, mono- or di-substituted isoxazolidin-2-yl, 4- and/or 5-, mono- and/or di-substituted isothiazolidin-2-yl wherein said R<sub>12</sub> substituents are independently H, halo, (C<sub>1</sub>-C<sub>5</sub>)alkyl, hydroxy, amino, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>5</sub>)alkylamino,  
30 formyl, oxo, hydroxyimino, (C<sub>1</sub>-C<sub>5</sub>)alkoxy, carboxy, carbamoyl, mono-N- or di-N,N-(C<sub>1</sub>-C<sub>5</sub>)alkylcarbamoyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxyimino, (C<sub>1</sub>-C<sub>4</sub>)alkoxymethoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxycarbonyl, carboxy(C<sub>1</sub>-C<sub>5</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>5</sub>)alkyl;


35 with the proviso that if R<sub>4</sub> is H, methyl, ethyl or n-propyl, R<sub>5</sub> is OH;

with the proviso that if  $R_5$  and  $R_7$  are H, then  $R_4$  is not H, methyl, ethyl, n-propyl, hydroxy( $C_1$ - $C_3$ )alkyl or ( $C_1$ - $C_3$ )alkoxy( $C_1$ - $C_3$ )alkyl and  $R_6$  is  $C(O)NR_8R_9$ ,  $C(O)R_{12}$  or ( $C_1$ - $C_4$ )alkoxycarbonyl;

5

and wherein Formula II is

10



Formula II

or the pharmaceutically acceptable salts or prodrugs thereof wherein the dotted line (---) is an optional bond wherein

A is  $-C(H)=$ ,  $-C((C_1-C_4)alkyl)=$ ,  $-C(halo)=$  or  $-N=$ , when the dotted line (---) is a bond, or A is methylene or  $-CH((C_1-C_4)alkyl)-$ , when the dotted line (---) is not a bond;

$R_1$ ,  $R_{10}$  or  $R_{11}$  are each independently H, halo, cyano, 4-, 6- or 7-nitro, ( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy, fluoromethyl, difluoromethyl or trifluoromethyl;

$R_2$  is H;

$R_3$  is H or ( $C_1$ - $C_5$ )alkyl;

$R_4$  is H, methyl, ethyl, n-propyl, hydroxy( $C_1$ - $C_3$ )alkyl, ( $C_1$ - $C_3$ )alkoxy( $C_1$ - $C_3$ )alkyl, phenyl( $C_1$ - $C_4$ )alkyl, phenylhydroxy( $C_1$ - $C_4$ )alkyl, (phenyl)(( $C_1$ - $C_4$ )-alkoxy)( $C_1$ - $C_4$ )alkyl, thien-2- or -3-yl( $C_1$ - $C_4$ )alkyl or fur-2- or -3-yl( $C_1$ - $C_4$ )alkyl wherein said  $R_4$  rings are mono-, di- or tri-substituted independently on carbon with H, halo, ( $C_1$ - $C_4$ )alkyl, ( $C_1$ - $C_4$ )alkoxy, trifluoromethyl, hydroxy, amino, cyano or 4,5-dihydro-1H-imidazol-2-yl; or

$R_4$  is pyrid-2-, -3- or -4-yl( $C_1$ - $C_4$ )alkyl, thiazol-2-, -4- or -5-yl( $C_1$ - $C_4$ )alkyl, imidazol-2-, -4-, or -5-yl( $C_1$ - $C_4$ )alkyl, pyrrol-2- or -3-yl( $C_1$ - $C_4$ )alkyl,

oxazol-2-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrazol-3-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, isoxazol-3-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, isothiazol-3-, -4- or -5-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyridazin-3- or -4-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrimidin-2-, -4-, -5- or  
5 -6-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, pyrazin-2- or -3-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl, 1,3,5-triazin-2-yl(C<sub>1</sub>-C<sub>4</sub>)alkyl or indol-2-(C<sub>1</sub>-C<sub>4</sub>)alkyl, wherein said preceding R<sub>4</sub> heterocycles are optionally mono- or di-substituted independently with halo, trifluoromethyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino,  
10 hydroxy or cyano and said substituents are bonded to carbon; or

R<sub>4</sub> is R<sub>15</sub>-carbonyloxymethyl, wherein said R<sub>15</sub> is phenyl, thiazolyl, imidazolyl, 1H-indolyl, furyl, pyrrolyl, oxazolyl, pyrazolyl, isoxazolyl, isothiazolyl,  
15 pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or 1,3,5-triazinyl and wherein said preceding R<sub>15</sub> rings are optionally mono- or di-substituted independently with halo, amino, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy or trifluoromethyl and said mono- or di-substituents are  
20 bonded to carbon;

R<sub>5</sub> is H, methyl, ethyl, n-propyl, hydroxymethyl or hydroxyethyl;

R<sub>6</sub> is carboxy, (C<sub>1</sub>-C<sub>8</sub>)alkoxycarbonyl, benzyloxycarbonyl, C(O)NR<sub>8</sub>R<sub>9</sub> or C(O)R<sub>12</sub>  
25 wherein R<sub>8</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>6</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>6</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkyl, hydroxy or (C<sub>1</sub>-C<sub>8</sub>)alkoxy; and

R<sub>9</sub> is H, cyclo(C<sub>3</sub>-C<sub>8</sub>)alkyl, cyclo(C<sub>3</sub>-C<sub>8</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkyl, cyclo(C<sub>4</sub>-C<sub>7</sub>)alkenyl,  
30 cyclo(C<sub>3</sub>-C<sub>7</sub>)alkyl(C<sub>1</sub>-C<sub>5</sub>)alkoxy, cyclo(C<sub>3</sub>-C<sub>7</sub>)alkyloxy, hydroxy, methylene-perfluorinated (C<sub>1</sub>-C<sub>8</sub>)alkyl, phenyl, or a heterocycle wherein said heterocycle is pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, pyridinyl, piperidinyl,  
35 morpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl, benzothiazolyl,

benzoxazolyl, benzimidazolyl, thiochromanyl or tetrahydrobenzothiazolyl wherein said heterocycle rings are carbon-nitrogen linked; or

$R_9$  is  $(C_1-C_6)$ alkyl or  $(C_1-C_8)$ alkoxy wherein said  
5  $(C_1-C_6)$ alkyl or  $(C_1-C_8)$ alkoxy is optionally monosubstituted with cyclo $(C_4-C_7)$ alken-1-yl, phenyl, thienyl, pyridyl, furyl, pyrrolyl, pyrrolidinyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, pyranyl, piperidinyl,  
10 morpholinyl, thiomorpholinyl, 1-oxothiomorpholinyl, 1,1-dioxothiomorpholinyl, pyridazinyl, pyrimidinyl, pyrazinyl, piperazinyl, 1,3,5-triazinyl or indolyl and wherein said  $(C_1-C_6)$ alkyl or  $(C_1-C_8)$ alkoxy are optionally additionally independently mono- or di-substituted with  
15 halo, hydroxy,  $(C_1-C_5)$ alkoxy, amino, mono-N- or di-N,N- $(C_1-C_5)$ alkylamino, cyano, carboxy, or  $(C_1-C_4)$ alkoxycarbonyl; and

wherein the  $R_9$  rings are optionally mono- or di-substituted independently on carbon with halo,  
20  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy, hydroxy, hydroxy $(C_1-C_4)$ alkyl, amino $(C_1-C_4)$ alkyl, mono-N- or di-N,N- $(C_1-C_4)$ alkylamino  $(C_1-C_4)$ alkyl,  $(C_1-C_4)$ alkoxy $(C_1-C_4)$ alkyl, amino, mono-N- or di-N,N- $(C_1-C_4)$ alkylamino, cyano, carboxy,  $(C_1-C_5)$ alkoxycarbonyl, carbamoyl, formyl or  
25 trifluoromethyl and said  $R_9$  rings may optionally be additionally mono- or di-substituted independently with  $(C_1-C_5)$ alkyl or halo;

with the proviso that no quaternized nitrogen on any  $R_9$  heterocycle is included;

30  $R_{12}$  is morpholino, thiomorpholino, 1-oxothiomorpholino, 1,1-dioxothiomorpholino, thiazolidin-3-yl, 1-oxothiazolidin-3-yl, 1,1-dioxothiazolidin-3-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, piperazin-4-yl,  
35 azetidin-1-yl, 1,2-oxazinan-2-yl, pyrazolidin-1-yl, isoxazolidin-2-yl, isothiazolidin-2-yl, 1,2-oxazetidin-2-yl, oxazolidin-3-yl,

3,4-dihydroisoquinolin-2-yl, 1,3-dihydroisoindol-2-yl,  
 3,4-dihydro-2H-quinol-1-yl, 2,3-dihydro-  
 benzo[1,4]oxazin-4-yl, 2,3-dihydro-benzo[1,4]-  
 thiazine-4-yl, 3,4-dihydro-2H-quinoxalin-1-yl,  
 5 3,4-dihydro-benzo[c][1,2]oxazin-1-yl, 1,4-dihydro-  
 benzo[d][1,2]oxazin-3-yl, 3,4-dihydro-benzo[e][1,2]-  
 oxazin-2-yl, 3H-benzo[d]isoxazol-2-yl,  
 3H-benzo[c]isoxazol-1-yl or azepan-1-yl,

wherein said  $R_{12}$  rings are optionally mono-, di-  
 10 or tri-substituted independently with halo,  $(C_1-C_5)$ alkyl,  
 $(C_1-C_5)$ alkoxy, hydroxy, amino, mono-N- or  
 di-N,N- $(C_1-C_5)$ alkylamino, formyl, carboxy, carbamoyl,  
 mono-N- or di-N,N- $(C_1-C_5)$ alkylcarbamoyl,  $(C_1-C_6)$ alkoxy  
 $(C_1-C_3)$ alkoxy,  $(C_1-C_5)$ alkoxycarbonyl, benzyloxycarbonyl,  
 15  $(C_1-C_5)$ alkoxycarbonyl  $(C_1-C_5)$ alkyl,  
 $(C_1-C_4)$ alkoxycarbonylamino, carboxy  $(C_1-C_5)$ alkyl,  
 carbamoyl  $(C_1-C_5)$ alkyl, mono-N- or  
 di-N,N- $(C_1-C_5)$ alkylcarbamoyl  $(C_1-C_5)$ alkyl,  
 hydroxy  $(C_1-C_5)$ alkyl,  $(C_1-C_4)$ alkoxy  $(C_1-C_4)$ alkyl,  
 20 amino  $(C_1-C_4)$ alkyl, mono-N- or  
 di-N,N- $(C_1-C_4)$ alkylamino  $(C_1-C_4)$ alkyl, oxo, hydroxyimino or  
 $(C_1-C_6)$ alkoxyimino and wherein no more than two  
 substituents are selected from oxo, hydroxyimino or  
 $(C_1-C_6)$ alkoxyimino and oxo, hydroxyimino or  
 25  $(C_1-C_6)$ alkoxyimino are on nonaromatic carbon; and

wherein said  $R_{12}$  rings are optionally  
 additionally mono- or di-substituted independently with  
 $(C_1-C_5)$ alkyl or halo;

with the proviso that when  $R_6$  is  
 30  $(C_1-C_5)$ alkoxycarbonyl or benzyloxycarbonyl then  $R_1$  is  
 5-halo, 5- $(C_1-C_4)$ alkyl or 5-cyano and  $R_4$  is  
 (phenyl) (hydroxy)  $(C_1-C_4)$ alkyl,  
 (phenyl)  $((C_1-C_4)$ alkoxy)  $(C_1-C_4)$ alkyl, hydroxymethyl or  
 Ar  $(C_1-C_2)$ alkyl, wherein Ar is thien-2- or -3-yl, fur-2- or  
 35 -3-yl or phenyl wherein said Ar is optionally mono- or  
 di-substituted independently with halo; with the provisos  
 that when  $R_4$  is benzyl and  $R_5$  is methyl,  $R_{12}$  is not

4-hydroxy-piperidin-1-yl or when  $R_4$  is benzyl and  $R_5$  is methyl  $R_6$  is not  $C(O)N(CH_3)_2$ ;

with the proviso that when  $R_1$  and  $R_{10}$  and  $R_{11}$  are H,  $R_4$  is not imidazol-4-ylmethyl, 2-phenylethyl or

5 2-hydroxy-2-phenylethyl;


with the proviso that when  $R_8$  and  $R_9$  are n-pentyl,  $R_1$  is 5-chloro, 5-bromo, 5-cyano, 5-( $C_1-C_5$ )alkyl, 5-( $C_1-C_5$ )alkoxy or trifluoromethyl;

with the proviso that when  $R_{12}$  is  
10 3,4-dihydroisoquinol-2-yl, said 3,4-dihydroisoquinol-2-yl is not substituted with carboxy-( $C_1-C_4$ )alkyl;

with the proviso that when  $R_8$  is H and  $R_9$  is ( $C_1-C_6$ )alkyl,  $R_9$  is not substituted with carboxy or ( $C_1-C_4$ )alkoxycarbonyl on the carbon which is attached to  
15 the nitrogen atom N of  $NHR_9$ ; and

with the proviso that when  $R_6$  is carboxy and  $R_1$ ,  $R_{10}$ ,  $R_{11}$  and  $R_5$  are all H, then  $R_4$  is not benzyl, H, (phenyl)(hydroxy)methyl, methyl, ethyl or n-propyl;

20 and wherein Formula III is



Formula III

30 or a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug wherein


$R^1$  is ( $C_1-C_4$ )alkyl, ( $C_1-C_7$ )cycloalkyl, phenyl or phenyl substituted with up to three ( $C_1-C_4$ )alkyl, ( $C_1-C_4$ )alkoxy or halogen;

35  $R^2$  is ( $C_1-C_4$ ) alkyl; and

$R^3$  is ( $C_3-C_7$ )cycloalkyl; phenyl; phenyl substituted at the para position with ( $C_1-C_4$ ) alkyl, halo,

hydroxy (C<sub>1</sub>-C<sub>4</sub>)alkyl or trifluoromethyl; phenyl substituted at the meta position with fluoro; or phenyl substituted at the ortho position with fluoro;

5 and wherein Formula IV is



Formula IV

15

a stereoisomer, pharmaceutically acceptable salt or prodrug thereof, or a pharmaceutically acceptable salt of the prodrug, wherein

20 Q is aryl, substituted aryl, heteroaryl, or substituted heteroaryl;

each Z and X are independently (C, CH or CH<sub>2</sub>), N, O or S; X<sup>1</sup> is NR<sup>a</sup>, -CH<sub>2</sub>-, O or S;

each - - - is independently a bond or is absent, provided that both - - - are not simultaneously bonds;


25 R<sup>1</sup> is hydrogen, halogen, -OC<sub>1</sub>-C<sub>8</sub>alkyl, -SC<sub>1</sub>-C<sub>8</sub>alkyl, -C<sub>1</sub>-C<sub>8</sub>alkyl, -CF<sub>3</sub>, -NH<sub>2</sub>, -NHC<sub>1</sub>-C<sub>8</sub>alkyl, -N(C<sub>1</sub>-C<sub>8</sub>alkyl)<sub>2</sub>, -NO<sub>2</sub>, -CN,

-CO<sub>2</sub>H, -CO<sub>2</sub>C<sub>1</sub>-C<sub>8</sub>alkyl, -C<sub>2</sub>-C<sub>8</sub>alkenyl, or -C<sub>2</sub>-C<sub>8</sub>alkynyl;


each R<sup>a</sup> and R<sup>b</sup> is independently hydrogen or -C<sub>1</sub>-C<sub>8</sub>alkyl;

30

35



- $R^2$  and  $R^3$  are independently hydrogen, halogen,  $-C_1-C_8$ alkyl,  $-CN$ ,  $-C\equiv C-Si(CH_3)_3$ ,  $-OC_1-C_8$ alkyl,  $-SC_1-C_8$ alkyl,  $-CF_3$ ,  $-NH_2$ ,  $-NHC_1-C_8$ alkyl,  $-N(C_1-C_8$ alkyl) $_2$ ,  
 5  $-NO_2$ ,  $-CO_2H$ ,  $-CO_2C_1-C_8$ alkyl,  $-C_2-C_8$ alkenyl, or  $-C_2-C_8$ alkynyl, or  $R^2$  and  $R^3$  together with the atoms on the ring to which they are attached form a five or six membered ring containing from 0 to 3 heteroatoms and from 0 to 2 double bonds;  
 10  $R^4$  is  $-C(=O)-A$ ;  
 A is  $-NR^dR^d$ ,  $-NR^aCH_2CH_2OR^a$ ,



- each  $R^d$  is independently hydrogen,  $C_1-C_8$ alkyl,  $C_1-C_8$ alkoxy, aryl, substituted aryl, heteroaryl, or substituted heteroaryl;  
 20 each  $R^c$  is independently hydrogen,  $-C(=O)OR^a$ ,  $-OR^a$ ,  $-SR^a$ , or  $-NR^aR^a$ ; and each n is independently 1-3.

- 25 3. A pharmaceutical composition comprising a glycogen phosphorylase inhibitor and a concentration-enhancing polymer, said glycogen phosphorylase inhibitor having a solubility in aqueous solution, in the absence of said concentration-enhancing polymer, of less than  
 30 1 mg/ml at any pH of from 1 to 8.

4. The composition of any one of claims 1-3 wherein said composition is a solid amorphous dispersion.

- 35 5. The composition of claim 4 wherein said dispersion is substantially homogeneous.



6. The composition of claim 4 wherein said glycogen phosphorylase inhibitor is almost completely amorphous.

5 7. The composition of any one of claims 1-3 wherein said composition is a simple physical mixture.

8. The composition of claim 7 wherein said mixture is substantially homogeneous.

10

9. The composition of claim 7 wherein said glycogen phosphorylase inhibitor is almost completely amorphous.

15 10. The composition of any one of claims 1-3 wherein said glycogen phosphorylase inhibitor is in a solid amorphous dispersion and only a portion of said concentration-enhancing polymer is present in said dispersion.

20

11. The composition of claim 1 wherein a portion of said glycogen phosphorylase inhibitor binds to one or more of the following residues of said glycogen phosphorylase enzyme in one or both subunits:

25

	parent secondary <u>structure</u>	<u>residue number</u>
		13-23
	helix $\alpha$ 1	24-37
30	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80
	strand $\beta$ 2	91-92
		93
35	helix $\alpha$ 3	94-102
		103

		121
	helix $\alpha$ 4	104-115
		116-117
	helix $\alpha$ 5	118-124
		125-128
5	strand $\beta$ 3	129-130
	strand $\beta$ 4	159-160
		161
	strand $\beta$ 4b	162-163
		164-166
10	strand $\beta$ 5	167-168
	strand $\beta$ 6	178
		179-190
	strand $\beta$ 7	191-192
		194, 197
15	strand $\beta$ 9	198-200
	strand $\beta$ 10	220-226
		228-232
		233-236
	strand $\beta$ 11	237-239, 241, 243-247
20		248-260
	helix $\alpha$ 7	261-276
	strand $\beta$ 11b	277-280

12. The composition of claim 1 wherein a  
 25 portion of said glycogen phosphorylase inhibitor binds to  
 a portion or all portions of the following residues of  
 said glycogen phosphorylase enzyme in one or both  
 subunits:

30	<u>residue number</u>
	33-39
	49-66
	94
	98
35	102
	125-126
	160

122

162  
 182-192  
 197  
 224-226  
 5 228-231  
 238-239  
 241  
 245  
 247

10

13. The composition of claim 1 wherein a  
 portion of said glycogen phosphorylase inhibitor binds to  
 a portion or all portions of the following residues of  
 said glycogen phosphorylase enzyme in one or both  
 15 subunits:

residue number  
 37-39  
 53  
 20 57  
 60  
 63-64  
 184-192  
 226  
 25 229

14. The composition of any one of claims 2-3  
 wherein a portion of said glycogen phosphorylase  
 inhibitor binds to a portion or all portions of the  
 30 following residues of a glycogen phosphorylase enzyme:

	<u>parent secondary structure</u>	<u>residue number</u>
		13-23
35	helix $\alpha$ 1	24-37
	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80

		123
	strand $\beta$ 1	81-86
		87-88
	strand $\beta$ 2	89-92
		93
5	helix $\alpha$ 3	94-102
		103
	helix $\alpha$ 4	104-115
		116-117
	helix $\alpha$ 5	118-124
10		125-128
	strand $\beta$ 3	129-131
		132-133
	helix $\alpha$ 6	134-150
		151-152
15	strand $\beta$ 4	153-160
		161
	strand $\beta$ 4b	162-163
		164-166
	strand $\beta$ 5	167-171
20		172-173
	strand $\beta$ 6	174-178
		179-190
	strand $\beta$ 7	191-192
		194, 197
25	strand $\beta$ 8	198-209
		210-211
	strand $\beta$ 9	212-216
	strand $\beta$ 10	219-226, 228-232
		233-236
30	strand $\beta$ 11	237-239, 241, 243-247
		248-260
	helix $\alpha$ 7	261-276
	strand $\beta$ 11b	277-281
	reverse turn	282-289
35	helix $\alpha$ 8	290-304

15. The composition of claim 14 wherein a portion of said glycogen phosphorylase inhibitor binds to a portion or all portions of the following residues of said glycogen phosphorylase enzyme in one or both subunits:

	parent secondary <u>structure</u>	<u>residue number</u>
		13-23
10	helix $\alpha$ 1	24-37
	turn	38-39, 43, 46-47
	helix $\alpha$ 2	48-66, 69-70, 73-74, 76-78
		79-80
	strand $\beta$ 2	91-92
15		93
	helix $\alpha$ 3	94-102
		103
	helix $\alpha$ 4	104-115
		116-117
20	helix $\alpha$ 5	118-124
		125-128
	strand $\beta$ 3	129-130
	strand $\beta$ 4	159-160
		161
25	strand $\beta$ 4b	162-163
		164-166
	strand $\beta$ 5	167-168
	strand $\beta$ 6	178
		179-190
30	strand $\beta$ 7	191-192
		194, 197
	strand $\beta$ 9	198-200
	strand $\beta$ 10	220-226
		228-232
35		233-236

125

strand $\beta$ 11	237-239, 241, 243-247
	248-260
helix $\alpha$ 7	261-276
strand $\beta$ 11b	277-280

5

16. The composition of claim 14 wherein a portion of said glycogen phosphorylase inhibitor binds to a portion or all portions of the following residues of said glycogen phosphorylase enzyme in one or both

10 subunits:

residue number

33-39

49-66

15

94

98

102

125-126

160

20

162

182-192

197

224-226

228-231

25

238-239

241

245

247

17. The composition of claim 14 wherein a portion of said glycogen phosphorylase inhibitor binds to a portion or all portions of the following residues of said glycogen phosphorylase enzyme in one or both subunits:

	<u>residue number</u>
	37-39
	53
10	57
	60
	63-64
	184-192
	226
15	229

18. The composition of claim 1 wherein said glycogen phosphorylase inhibitor has the structure of Formula I defined in claim 2.

20

19. The composition of claim 2 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-chloro-1H-indole-2-carboxylic acid [(1S)-(R)-hydroxy-dimethylcarbamoylmethyl]-2-phenyl-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-(R)-hydroxy-methoxy-methylcarbamoylmethyl]-2-phenyl-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3S)-hydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4R)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, and 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-morpholin-4-yl-3-oxo-propyl]-amide.

20. The composition of claim 18 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-dimethylcarbamoylmethyl)-2-phenyl-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-((R)-hydroxy-methoxy-methylcarbamoylmethyl)-2-phenyl-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3S)-hydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4R)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide, and 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-morpholin-4-yl-3-oxo-propyl]-amide.

21. The composition of claim 1 wherein said glycogen phosphorylase inhibitor has the structure of Formula II defined in claim 2.

20

22. The composition of claim 2 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-chloro-1H-indole-2-carboxylic acid [2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide, and 5-chloro-1H-indole-2-carboxylic acid [2-(1-oxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide.



23. The composition of claim 21 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-chloro-1H-indole-2-carboxylic acid [2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3S,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-((3R,4S)-3,4-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-(4-fluoro-benzyl)-2-(4-hydroxy-piperidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-2-(3-hydroxy-azetidin-1-yl)-2-oxo-ethyl]-amide, 5-chloro-1H-indole-2-carboxylic acid [2-(1,1-dioxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide, and 5-chloro-1H-indole-2-carboxylic acid [2-(1-oxo-thiazolidin-3-yl)-2-oxo-ethyl]-amide.

24. The composition of claim 1 wherein said glycogen phosphorylase inhibitor has the structure of Formula III as defined in claim 2.

25. The composition of claim 2 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-acetyl-1-ethyl-2,3-dihydro-2-oxo-N-[3-[(phenylamino)carbonyl]phenyl]-1H-Indole-3-carboxamide, 5-acetyl-N-[3-[(cyclohexylamino)carbonyl]phenyl]-1-ethyl-2,3-dihydro-2-oxo-1H-Indole-3-carboxamide, and 5-acetyl-N-[3-[(4-bromophenyl)amino]carbonyl]phenyl]-2,3-dihydro-1-methyl-2-oxo-1H-Indole-3-carboxamide.

26. The composition of claim 24 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 5-acetyl-1-ethyl-2,3-dihydro-2-oxo-N-[3-[(phenylamino)carbonyl]phenyl]-1H-Indole-3-carboxamide, 5-acetyl-N-[3-[(cyclohexylamino)carbonyl]phenyl]-1-ethyl-2,3-dihydro-2-

oxo-1H-Indole-3-carboxamide, and 5-acetyl-N-[3-[(4-bromophenyl)amino]carbonyl]phenyl]-2,3-dihydro-1-methyl-2-oxo-1H-Indole-3-carboxamide.

5                   27. The composition of claim 1 wherein said glycogen phosphorylase inhibitor has the structure of Formula IV as defined in claim 2.

10                   28. The composition of claim 2 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 2-Chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, and 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-(2R)-  
15                   hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide.

20                   29. The composition of claim 27 wherein said glycogen phosphorylase inhibitor is selected from the group consisting of 2-Chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-2-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-2-oxo-ethyl]-amide, and 2-chloro-6H-thieno[2,3-b]pyrrole-5-carboxylic acid [(1S)-benzyl-(2R)-  
25                   hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxo-propyl]-amide.

30                   30. The composition of any one of claims 1 and 2 wherein said glycogen phosphorylase inhibitor has a solubility in aqueous solution in the absence of said  
30                   concentration-enhancing polymer of less than 1 mg/ml at any pH of from 1 to 8.

35                   31. The composition of claim 30 wherein said glycogen phosphorylase inhibitor has an aqueous solubility of less than 0.5 mg/ml.

32. The composition of claim 3 wherein said glycogen phosphorylase inhibitor has an aqueous solubility of less than 0.5 mg/ml.

5           33. The composition of claim 31 wherein said solubility is less than 0.1 mg/mL.

34. The composition of claim 32 wherein said solubility is less than 0.1 mg/mL.

10

35. The composition of any one of claims 1-3 wherein said glycogen phosphorylase inhibitor has a dose-to-aqueous-solubility ratio of at least 10 ml.

15

36. The composition of claim 35 wherein said dose-to-aqueous solubility ratio is at least 100 ml.

37. The composition of claim 36 wherein said dose-to-aqueous solubility ratio is at least 400 ml.

20

38. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer comprises a blend of polymers.

25

39. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer has at least one hydrophobic portion and at least one hydrophilic portion.

30

40. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer is an ionizable polymer.

35

41. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer is selected from the group consisting of ionizable cellulosic polymers, nonionizable cellulosic polymers, and vinyl

polymers and copolymers having substituents selected from the group consisting of hydroxyl, alkylacyloxy, and cyclicamido.

5                   42. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer is a cellulosic polymer.

10                   43. The composition of claim 42 wherein said concentration-enhancing polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and  
15                   hydroxyethyl ethyl cellulose.

                  44. The composition of claim 42 wherein said concentration-enhancing polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose  
20                   acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, hydroxypropyl  
25                   methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose  
30                   butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate

trimellitate, hydroxypropyl methyl cellulose acetate  
trimellitate, hydroxypropyl cellulose acetate  
trimellitate succinate, cellulose propionate  
trimellitate, cellulose butyrate trimellitate, cellulose  
5 acetate terephthalate, cellulose acetate isophthalate,  
cellulose acetate pyridinedicarboxylate, salicylic acid  
cellulose acetate, hydroxypropyl salicylic acid cellulose  
acetate, ethylbenzoic acid cellulose acetate,  
hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl  
10 phthalic acid cellulose acetate, ethyl nicotinic acid  
cellulose acetate, and ethyl picolinic acid cellulose  
acetate.

45. The composition of claim 42 wherein said  
15 concentration-enhancing polymer is selected from the  
group consisting of cellulose acetate phthalate, methyl  
cellulose acetate phthalate, ethyl cellulose acetate  
phthalate, hydroxypropyl cellulose acetate phthalate,  
hydroxypropyl methyl cellulose phthalate, hydroxypropyl  
20 methyl cellulose acetate phthalate, hydroxypropyl  
cellulose acetate phthalate succinate, cellulose  
propionate phthalate, hydroxypropyl cellulose butyrate  
phthalate, cellulose acetate trimellitate, methyl  
cellulose acetate trimellitate, ethyl cellulose acetate  
25 trimellitate, hydroxypropyl cellulose acetate  
trimellitate, hydroxypropyl methyl cellulose acetate  
trimellitate, hydroxypropyl cellulose acetate  
trimellitate succinate, cellulose propionate  
trimellitate, cellulose butyrate trimellitate, cellulose  
30 acetate terephthalate, cellulose acetate isophthalate,  
cellulose acetate pyridinedicarboxylate, salicylic acid  
cellulose acetate, hydroxypropyl salicylic acid cellulose  
acetate, ethylbenzoic acid cellulose acetate,  
hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl  
35 phthalic acid cellulose acetate, ethyl nicotinic acid  
cellulose acetate, and ethyl picolinic acid cellulose  
acetate.

46. The composition of claim 42 wherein said concentration-enhancing polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate succinate, cellulose acetate phthalate, hydroxypropyl methyl cellulose phthalate, methyl cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate and cellulose acetate isophthalate.

47. The composition of claim 46 wherein said concentration-enhancing polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, and cellulose acetate trimellitate.

48. The composition of any one of claims 1-3 wherein said concentration-enhancing polymer is present in an amount sufficient to permit said composition to provide a maximum concentration of said glycogen phosphorylase inhibitor in a use environment that is at least 1.25-fold that of a control composition comprising an equivalent quantity of said glycogen phosphorylase inhibitor and free from said concentration-enhancing polymer.

49. The composition of claim 48 wherein said maximum concentration of said glycogen phosphorylase inhibitor in said use environment is at least 2-fold that of said control composition.

50. The composition of any one of claims 1-3 wherein said composition provides in an aqueous use environment an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into the use environment and about

270 minutes following introduction to the use environment that is at least 1.25-fold that of a control composition comprising an equivalent quantity of said glycogen phosphorylase inhibitor and free from said concentration-enhancing polymer.

51. The composition of any one of claims 1-3 wherein said composition provides a relative bioavailability that is at least 1.25 relative to a control composition comprising an equivalent quantity of said glycogen phosphorylase inhibitor and free from said concentration-enhancing polymer.

52. The composition of claim 48 wherein said use environment is *in vitro*.

53. The composition of claim 48 wherein said use environment is *in vivo*.

54. The composition of claim 53 wherein said use environment is the gastrointestinal tract of an animal.

55. The composition of claim 54 wherein said animal is a human.

56. The composition of claim 50 wherein said use environment is *in vitro*.

57. The composition of claim 50 wherein said use environment is *in vivo*.

58. The composition of claim 57 wherein said use environment is the gastrointestinal tract of an animal.

59. The composition of claim 58 wherein said animal is a human.

60. The composition of claim 4 wherein said dispersion is formed by solvent processing.

61. The composition of claim 60 wherein said solvent processing is spray-drying.

62. A method of treating diabetes, the method comprising the step of administering to a patient having diabetes a therapeutically effective amount of a composition of any one of claims 1-3.

63. The method of claim 62 wherein the diabetes is non-insulin dependent diabetes mellitus (Type 2).

64. The method of claim 62 wherein the diabetes is insulin dependent diabetes mellitus (Type 1).

65. A method of treating or presenting an indication selected from the group consisting of atherosclerosis, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, cataracts, hypercholesterolemia, hypertriglyceridemia, hyperlipidemia, hyperglycemia, hypertension, tissue ischemia, myocardial ischemia, insulin resistance, bacterial infection, diabetic cardiomyopathy and tumor growth, the method comprising the step of administering to a patient a therapeutically effective amount of a composition of any one of claims 1-3.



66. A method of inhibiting glycogen phosphorylase, the method comprising the step of administering to a patient in need of glycogen phosphorylase inhibition, a glycogen phosphorylase inhibiting amount of a composition of any one of claims 1-3.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 01/00394

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K9/14

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 901 786 A (PFIZER PROD INC) 17 March 1999 (1999-03-17) page 3, line 15 -page 5, line 30 page 11 -page 12; example 1 claims 1,28-30 ---	1-66
A	WO 96 39385 A (TREADWAY JUDITH L ;HULIN BERNARD (US); HOOVER DENNIS J (US); PFIZE) 12 December 1996 (1996-12-12) cited in the application the whole document ---	1-66
A	WO 96 39384 A (TREADWAY JUDITH L ;HULIN BERNARD (US); HOOVER DENNIS J (US); PFIZE) 12 December 1996 (1996-12-12) cited in the application the whole document ---	1-66
-/-		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

16 August 2001

Date of mailing of the international search report

23/08/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Muller, S

# INTERNATIONAL SEARCH REPORT

International Applic.  
PCT/IB 01/00394

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 5 998 463 A (HULIN BERNARD ET AL)  7 December 1999 (1999-12-07)  cited in the application  the whole document  -----</p>	1-66

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-41,48-66

Present claims 1,4-18,20,21,23,24,26,27,29,30,31,33,35-41,48-66 relate to a composition defined by reference to a desirable characteristic, namely a portion of the glycogen phosphorylase inhibitor that binds to a portion or all portions of the following residues of a glycogen phosphorylase enzyme. The claims cover all compositions having this characteristic, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such compositions. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the composition by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Present claims 2,4-10,14-31,33,35-41,48-66 relate to an extremely large number of possible compositions. In fact, the claims contain so many options, variables, possible permutations and provisos that a lack of conciseness within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible.

Present claims 3-10,14-17,30-41,48-66 relate to a composition C comprising a concentration-enhancing polymer (CEP) and a glycogen phosphorylase inhibitor (GPI) defined by reference to the following parameter:

P1: said GPI having a solubility in aqueous solution, in the absence of said-concentration-enhancing polymer, of less than 1mg/ml at any pH of from 1 to 8.

Present claims 35-41 relate to said composition C defined by reference to the following parameter:

P2: wherein said GPI has a dose-to-aqueous-solubility ratio of at least 10ml.

Present claims 48,49,52-55 relate to said composition C defined by reference to the following parameter:

P3: wherein said CEP is present in an amount sufficient to permit said composition to provide a maximum concentration of said GPI in a defined use environment.

The use of these parameters in the present context is considered to lead to a lack of clarity within the meaning of Article 6 PCT. It is impossible to compare the parameters the applicant has chosen to employ with what is set out in the prior art. The lack of clarity is such as to render a meaningful complete search impossible.

Consequently, the search has been carried out for those parts of the application which do appear to be clear and concise, namely for compositions comprising the glycogen phosphorylase inhibitor compounds recited in the examples, namely "drug1" as defined in example1, "drug2" as defined in examples 12-17, and "drug3" as defined in examples 20-25

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

and the concentration-enhancing polymers recited in claims 42-47.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 01/00394

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0901786 A	17-03-1999	BR 9803144 A	11-01-2000
		CN 1207896 A	17-02-1999
		JP 2984661 B	29-11-1999
		JP 11116502 A	27-04-1999
WO 9639385 A	12-12-1996	CA 2223625 A	12-12-1996
		AP 624 A	19-12-1997
		AU 700887 B	14-01-1999
		AU 5475396 A	19-12-1996
		BG 62566 B	29-02-2000
		BG 100635 A	30-09-1997
		BR 9602626 A	01-09-1998
		CZ 9601627 A	11-12-1996
		EP 0832066 A	01-04-1998
		FI 974437 A	05-12-1997
		HR 960266 A	31-08-1997
		HU 9601285 A	28-09-1998
		JP 11500445 T	12-01-1999
		JP 3068200 B	24-07-2000
		LV 11614 A	20-12-1996
		LV 11614 B	20-04-1997
		NO 962322 A	09-12-1996
		PL 314603 A	09-12-1996
		SG 45481 A	16-01-1998
		SI 9600163 A	28-02-1997
		SK 72096 A	05-11-1997
		TR 970184 A	21-03-1997
WO 9639384 A	12-12-1996	CA 2224062 A	12-12-1996
		AP 623 A	19-12-1997
		AU 701465 B	28-01-1999
		AU 5462696 A	19-12-1996
		BG 100547 A	31-12-1996
		BR 9602542 A	27-10-1998
		CN 1142492 A	12-02-1997
		CN 1140709 A	22-01-1997
		CZ 9601573 A	12-03-1997
		EP 0832065 A	01-04-1998
		FI 974436 A	27-01-1998
		HR 960244 A	31-12-1997
		HU 9601475 A	28-09-1998
		JP 10511687 T	10-11-1998
		KR 191992 B	15-06-1999
		LV 11613 A	20-12-1996
		LV 11613 B	20-04-1997
		NO 961664 A	09-12-1996
		NO 990405 A	28-01-1999
		NZ 286460 A	24-09-1998
		PL 314561 A	09-12-1996
		RU 2143424 C	27-12-1999
		SG 44947 A	19-12-1997
		SI 9600177 A	28-02-1997
		SK 69996 A	06-08-1997
		TR 961048 A	21-12-1996
		US 6107329 A	22-08-2000
US 5998463 A	07-12-1999	NONE	